



Patent

Op 1638
Bk

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

Applicant(s): Schmülling et al.

Examiner: S. Baum

Serial No.: 10/014,101

Group: Art Unit 1638

Filed: December 10, 2001

Docket: 1187-26

For: **METHOD FOR ENHANCING
MODIFYING PLANT MORPHOLOGY,
BIOCHEMISTRY AND PHYSIOLOGY**

Dated: November 17, 2004


Commissioner for Patents
P.O. Box 1450
Alexandria, VA 22313-1450

TRANSMITTAL OF PRIORITY DOCUMENT

Sir:

Applicants in the above-identified application hereby claim the right of priority in connection with Title 35 U.S.C. §119 and in support thereof, herewith submit a certified copy of European Patent Application Number EP 00870132.8 filed June 16, 2000.

Respectfully submitted,



Ann R. Pokalsky
Reg. No. 34,697
Attorney for Applicants

DILWORTH & BARRESE
333 Earle Ovington Blvd.
Uniondale, NY 11553
(516) 228-8484
ARP:bg

CERTIFICATE OF MAILING UNDER 37 C.F.R. §1.8(a)

I hereby certify that this correspondence is being deposited with the United States Postal Service as first class mail, postpaid in an envelope, addressed to the: U.S. Patent and Trademark Office, P.O. Box 1450, Alexandria, VA 22313-1450 on November 17, 2004.

Dated: November 17, 2004


Bridget Griffin

THIS PAGE BLANK (USPTO)



**Europäisches
Patentamt**

**European
Patent Office**

**Office européen
des brevets**

Bescheinigung

Certificate

Attestation

Die angehefteten Unterlagen stimmen mit der ursprünglich eingereichten Fassung der auf dem nächsten Blatt bezeichneten europäischen Patentanmeldung überein.

The attached documents are exact copies of the European patent application described on the following page, as originally filed.

Les documents fixés à cette attestation sont conformes à la version initialement déposée de la demande de brevet européen spécifiée à la page suivante.

Patentanmeldung Nr. Patent application No. Demande de brevet n°

00870132.8

Der Präsident des Europäischen Patentamts;
Im Auftrag

For the President of the European Patent Office

Le Président de l'Office européen des brevets
p.o.

R C van Dijk

DEN HAAG, DEN
THE HAGUE, 28/09/04
LA HAYE, LE

THIS PAGE BLANK (USPTO)



Anmeldung Nr:
Application no.: 00870132.8
Demande no:

Anmeldetag:
Date of filing: 16.06.00
Date de dépôt:

Anmelder/Applicant(s)/Demandeur(s):

CropDesign N.V.
Technologiepark 3
9052 Zwijnaarde-Gent
BELGIQUE

Bezeichnung der Erfindung/Title of the invention/Titre de l'invention:
(Falls die Bezeichnung der Erfindung nicht angegeben ist, siehe Beschreibung.
If no title is shown please refer to the description.
Si aucun titre n'est indiqué se référer à la description.)

Method for modifying plant morphology, biochemistry and physiology.

In Anspruch genommene Priorität(en) / Priority(ies) claimed /Priorité(s)
revendiquée(s)
Staat/Tag/Aktenzeichen/State/Date/File no./Pays/Date/Numéro de dépôt:

/00.00.00/

Internationale Patentklassifikation/International Patent Classification/
Classification internationale des brevets:

C12N15/00

Am Anmeldetag benannte Vertragstaaten/Contracting states designated at date of
filing/Etats contractants désignées lors du dépôt:

AT BE CH CY DE DK ES FI FR GB GR IE IT LI LU MC NL PT SE

THIS PAGE BLANK (USPTO)

METHOD FOR MODIFYING PLANT MORPHOLOGY, BIOCHEMISTRY AND PHYSIOLOGY

FIELD OF THE INVENTION

5

The present invention generally relates to a method for modifying plant morphological, biochemical and physiological properties or characteristics, such as one or more developmental processes and/or environmental adaptive processes, including but not limited to the modification of initiation or stimulation
10 or enhancement of root growth, and/or adventitious root formation, and/or lateral root formation, and/or root geotropism, and/or shoot growth, and/or apical dominance, and/or branching, and/or timing of senescence, and/or timing of flowering, and/or flower formation, and/or seed development, and/or seed yield, said method comprising expressing a cytokinin degradation control protein, in
15 particular cytokinin oxidase, in the plant, operably under the control of a regulatable promoter sequence such as a cell-specific promoter, tissue-specific promoter, or organ-specific promoter sequence. Preferably, the characteristics modified by the present invention are cytokinin-mediated and/or auxin-mediated characteristics. The present invention extends to genetic constructs which are
20 useful for performing the inventive method and to transgenic plants produced therewith having altered morphological and/or biochemical and/or physiological properties compared to their otherwise isogenic counterparts.

BACKGROUND OF THE INVENTION

25

Roots are an important organ of higher plants. Their main functions are anchoring of the plant in the soil and uptake of water and nutrients (N-nutrition, minerals, etc.). Thus, root growth has a direct or indirect influence on growth and yield of aerial organs, particularly under conditions of nutrient limitation. Roots
30 are also relevant for the production of secondary plant products, such as defense compounds and plant hormones.

Roots are also storage organs in a number of important staple crops. Sugar beet is the most important plant for sugar production in Europe (260 Mill t/year; 38 % of world production). Manioc (cassava), yams and sweet potato (batate) are important starch producers (app. 150 Mill t/year each). Their content in starch
5 can be twice as high as that of potato. Roots are also the relevant organ for consumption in a number of vegetables (e.g. carrots, radish), herbs (e.g. ginger, kukuma) and medicinal plants (e.g. ginseng). In addition, some of the secondary plant products found in roots are of economic importance for the chemical industry. An example is yams, which contain basic molecules for the synthesis of
10 steroid hormones.

Moreover, improved root growth of crop plants will also enhance competitiveness with weedy plants and will improve growth in arid areas, by increasing water accessibility and uptake.

Improved root growth is also relevant for ecological purposes, such as
15 bioremediation and prevention/arrest of soil erosion.

Root architecture is an area that has remained largely unexplored through classical breeding, because of difficulties with assessing this trait in the field. Thus, biotechnology could have significant impact on the improvement of this trait, because it does not rely on large-scale screenings in the field. Rather,
20 biotechnological approaches require a basic understanding of the molecular components that determine a specific characteristic of the plant. Today, this knowledge is only fragmentary, and as a consequence, biotechnology was so far unable to realize a break-through in this area.

A well-established regulator of root growth is auxin. Application of indole-3-acetic acid (IAA) to growing plants stimulates lateral root development and lateral root elongation (Torrey, Am J Bot 37: 257-264, 1950; Blakely *et al.*, Bot Gaz 143: 341-352, 1982; Muday and Haworth, Plant Physiol Biochem 32: 193-203, 1994). Roots exposed to a range of concentrations of IAA initiated increasing numbers of lateral roots (Kerk *et al.*, Plant Physiol, 122: 925-932, 2000). Furthermore,
30 when roots that had produced laterals in response to a particular concentration of exogenous auxin were subsequently exposed to a higher concentration of IAA,

numerous supernumerary lateral roots spaced between existing ones were formed (Kerk *et al.*, Plant Physiol, 122: 925-932, 2000). Conversely, growth of roots on agar containing auxin-transport inhibitors, including NPA, decreases the number of lateral roots (Muday and Haworth, Plant Physiol Biochem 32: 193-203, 5 1994).

Arabidopsis mutants containing increased levels of endogenous IAA have been isolated (Boerjan *et al.*, Plant Cell 7: 1405-141, 1995; Celenza *et al.*, Gene Dev 9: 2131-2142, 1995; King *et al.*, Plant Cell 7: 2023-2037, 1995; Lehman *et al.*, Cell 85: 183-194, 1996). They are now known to be alleles of a single locus 10 located on chromosome 2. These mutant seedlings have excess adventitious and lateral roots, which is in accordance with the above-described effects of external auxin application.

The stimulatory effect of auxins on adventitious and lateral root formation suggests that overproduction of auxins in transgenic plants is a valid strategy for 15 increasing root growth. Yet, it is also questionable whether this would yield a commercial product with improved characteristics. Apart from its stimulatory effect on adventitious and lateral root formation, auxin overproduction triggers other effects, such as reduction in leaf number, abnormal leaf morphology (narrow, curled leaves), aborted inflorescences, increased apical dominance, 20 adventitious root formation on the stem, most of which are undesirable from an agronomic perspective (Klee *et al.*, Genes Devel 1: 86-96, 1987; Kares *et al.*, Plant Mol Biol 15: 225-236, 1990). Therefore, the major problem with approaches that rely on increased auxin synthesis is a problem of containment, namely to confine the effects of auxin to the root. This problem of containment is not likely 25 overcome by using tissue-specific promoters: auxins are transported in the plant and their action is consequently not confined to the site of synthesis. Another issue is whether auxins will always enhance the total root biomass. For agar-grown plants, it has been noticed that increasing concentrations progressively stimulated lateral root formation but concurrently inhibited the outgrowth of these 30 roots (Kerk *et al.*, Plant Physiol, 122: 925-932, 2000).

WO 99/06571 discloses a *Zea mays* cytokinin oxidase gene and the protein encoding the same as well as transgenic plants comprising the same.

The above-mentioned problems related to containment of auxin effects and to maintenance of root outgrowth are solved by the embodiments in the patent
5 claim.

SUMMARY OF THE INVENTION

The present invention relates to a genetic construct comprising a gene encoding
10 a protein with cytokinin oxidase activity from *Arabidopsis thaliana*. This gene is expressed under control of a regulated promoter. This promoter may be regulated by endogenous tissue-specific or environment-specific factors or, alternatively, it may be induced by application of specific chemicals.

The present invention also relates to a cell or plant containing the genetic
15 construct.

The present invention also relates to a method to modify root architecture and biomass by expression of a cytokinin oxidase gene under control of a promoter that is specific to the root or to certain tissues or cell types of the root.

20 DETAILED DESCRIPTION OF THE INVENTION

To by-pass above-mentioned problems associated with increasing auxin biosynthesis, it was decided to follow an alternative approach. We reasoned that down-regulation of auxin antagonists could evoke similar or even superior effects
25 on root growth as compared to increasing auxin levels. Hormone actions and interactions are extremely complex, but we hypothesized that cytokinins could function as auxin antagonists with respect to root growth. Hormone studies on plant tissue cultures have shown that the ratio of auxin versus cytokinin is more important for organogenesis than the absolute levels of each of these hormones,
30 which indeed indicates that these hormones function as antagonists – at least in certain biological processes. Furthermore, lateral root formation is inhibited by

exogenous application of cytokinins. Interestingly, also root elongation is negatively affected by cytokinin treatment, which suggests that cytokinins control both root branching and root outgrowth. .

Together, current literature data indicate that increasing cytokinin levels
5 negatively affects root growth, but the mechanisms underlying this process are not understood. The sites of cytokinin synthesis in the plant are root tips and young tissues of the shoot. Endogenous concentrations of cytokinins are in the nM range. However, as their quantification is difficult, rather large tissue amounts need to be extracted and actual local concentrations are not known. Also the
10 subcellular compartmentation of cytokinins is not known. It is generally thought that the free base and ribosides are localized in the cytoplasm and nucleus, while glucosides are localized in the vacuole. There exist also different cytokinins with slightly different chemical structure. As a consequence, it is not known whether the effects of exogenous cytokinins should be ascribed to a raise in total
15 cytokinin concentration or rather to the competing out of other forms of plant-borne cytokinins (which differ either in structure, cellular or subcellular location) for receptors, translocators, transporters, modifying enzymes ...

In order to test the hypothesis that cytokinin levels in the root indeed exceed the level optimal for root growth, novel genes encoding cytokinin oxidases (which are
20 cytokinin metabolizing enzymes) were cloned from *Arabidopsis thaliana* (designated *AtCKX*) and were subsequently expressed under a strong constitutive promoter in transgenic tobacco and *Arabidopsis*. Transformants showing *AtCKX* mRNA expression and increased cytokinin oxidase activity also manifested enhanced formation and growth of roots. Negative effects on shoot
25 growth were also observed. The latter is in accordance with the constitutive expression of the cytokinin oxidase gene in these plants, illustrating the importance of confined expression of the cytokinin oxidase gene for general plant growth properties. Containment of cytokinin oxidase activity can be achieved by using cell-, tissue- or organ-specific promoters, since cytokinin degradation is a
30 process limited to the cells that express the CKX protein, this in contrast to approaches relying on hormone synthesis, as was explained above.

The observed negative effects of cytokinin oxidase expression on shoot growth demonstrate that cytokinin oxidases are interesting targets for the design of or screening for growth-promoting chemicals. Such chemicals should inhibit cytokinin oxidase activity, should preferably not be transported to the root and should be rapidly degraded in soil, so that application of these chemicals will not inhibit root growth. Cytokinins also delay leaf senescence, which means that positive effects will include both growth and maintenance of photosynthetic tissues. . In addition, the observation that cytokinins delay senescence, enhance greening (chlorophyll content) of leaves and reduce shoot apical dominance shows that strategies based on suppressing CKX activity (such as antisense, ribozyme, and cosuppression technology) in the aerial parts of the plant could result in delayed senescence, enhanced leaf greening and increased branching. Similarly, the observed positive effects of cytokinin oxidase expression on root growth demonstrate that cytokinin oxidases are interesting targets for the design of or screening for herbicides. Such herbicides should inhibit cytokinin oxidase activity, should preferably not be transported to the shoot, and should be soluble and relatively stable in a solvent that can be administered to the root through the soil.

These effects of cytokinin oxidase overexpression on plant development and architecture were hitherto unknown and, as a consequence, the presented invention and its embodiments could not be envisaged.

Preferable embodiments of the invention relate to the positive effect of cytokinin oxidase expression on plant growth and architecture, and in particular on root growth and architecture. The cytokinin oxidase gene family contains at least six members in *Arabidopsis* (see examples below) and we have shown that there are quantitative differences in the effects achieved with some of these genes in transgenic plants. It is anticipated that functional homologs of the described *Arabidopsis* cytokinin oxidases can be isolated from other organisms, given the evidence for the presence of cytokinin oxidase activity in many green plants (Hare and van Staden, *Physiol Plant* 91:128-136, 1994; Jones and Schreiber, *Plant Growth Reg* 23:123-134, 1997), as well as in other organisms (Armstrong,

in Cytokinins: Chemistry, Activity and Function. Eds Mok and Mok, CRC Press, pp139-154, 1994). Therefore, the sequence of the cytokinin oxidase, functional in the invention, need not to be identical to those described herein. It is envisaged that other genes with cytokinin oxidase activity or with any other cytokinin metabolizing activity (see Zažímalová *et al.*, Biochemistry and Molecular Biology of Plant Hormones, Hooykaas, Hall and Libbenga (Eds.), Elsevier Science, pp141-160, 1997) can also be used for the purpose of this invention. Similarly, genes encoding proteins that would increase endogenous cytokinin metabolizing activity can also be used for the purpose of this invention.

10 According to a first embodiment, the present invention relates to a method for stimulating root growth and/or enhancing the formation of lateral and adventitious roots and/or altering root geotropism comprising expression of a plant cytokinin oxidase or comprising expression of another protein that reduces the level of active cytokinins in plants or plant parts.

15 In the context of the present invention it should be understood that the term "expression" relates to an "enhanced and/or ectopic expression" of a plant cytokinin oxidase or any other protein that reduces the level of active cytokins in plants. It should be clear that herewith an enhanced expression of the plant cytokinin oxidase as well as "de novo" expression of plant cytokinin oxidases or
20 of said other proteins is meant.

Alternatively, said other protein enhances the cytokinin metabolizing activity of a plant cytokinin oxidase.

According to a further embodiment, the present invention relates to a method for stimulating root growth and/or enhancing the formation of lateral and adventitious
25 roots and/or altering root geotropism and/or increasing yield and/or enhancing early vigor and/or modifying root/shoot ratio and/or improving lodging and/or increasing drought tolerance and/or promoting in vitro propagation of explants, comprising expression of a plant cytokinin oxidase or comprising expression of another protein that reduces the level of active cytokinins in plants or plant parts.

30 According to another embodiment, the present invention relates to an isolated nucleic acid encoding a novel plant protein having cytokinin oxidase activity or

encoding an immunologically active and/or functional fragment of such a protein selected from the group consisting of:

- (a) nucleic acid sequences comprising a DNA sequence as given in any of SEQ ID NOs 1, 3, 5, 7, 9 or 11, or the complement thereof,
- 5 (b) nucleic acid sequences comprising the RNA sequences corresponding to any of SEQ ID NOs 1, 3, 5, 7, 9 or 11, or the complement thereof,
- (c) nucleic acid sequences hybridizing to the nucleotide sequence as defined in (a) or (b) under stringent hybridization conditions,
- 10 (d) nucleic acid sequences encoding a protein with an amino acid sequence which is at least 35% similar, preferably 36%, 37%, 38%, 39% similar, more preferably 40%, 41%, 42%, 43%, 44% similar, most preferably 45%, 50%, 60%, 70%, 80% 90% or 95% similar to the amino acid sequence as given in SEQ ID NO 2,
- 15 (e) nucleic acid sequences encoding a protein with an amino acid sequence which is at least 35% similar, preferably 36%, 37%, 38%, 39% similar, more preferably 40%, 41%, 42%, 43%, 44% similar, most preferably 45%, 50%, 60%, 70%, 80% 90% or 95% similar to the amino acid sequence as given in SEQ ID NO 4,
- 20 (f) nucleic acid sequences encoding a protein with an amino acid sequence which is at least 35% similar, preferably 36%, 37%, 38%, 39% similar, more preferably 40%, 41%, 42%, 43%, 44% similar, most preferably 45% 50%, 60%, 70%, 80% 90% or 95% similar to the amino acid sequence as given in SEQ ID NO 6,
- 25 (g) nucleic acid sequences encoding a protein with an amino acid sequence which is at least 35% similar, preferably 36%, 37%, 38%, 39% similar, more preferably 40%, 41%, 42%, 43%, 44% similar, most preferably 45% 50%, 60%, 70%, 80% 90% or 95% similar to the amino acid sequence as given in SEQ ID NO 8,
- 30 (h) nucleic acid sequences encoding a protein with an amino acid sequence which is at least 35% similar, preferably 36%, 37%, 38%, 39% similar, more preferably 40%, 41%, 42%, 43%, 44% similar, most preferably 45%

50%, 60%, 70%, 80% 90% or 95% similar to the amino acid sequence as given in SEQ ID NO 10,

- 5 (i) nucleic acid sequences encoding a protein with an amino acid sequence which is at least 35% similar, preferably 36%, 37%, 38%, 39% similar, more preferably 40%, 41%, 42%, 43%, 44% similar, most preferably 45% 50%, 60%, 70%, 80% 90% or 95% similar to the amino acid sequence as given in SEQ ID NO 12,
- (j) nucleic acid sequences encoding a protein comprising the amino acid sequence as given in any of SEQ ID NOs 2, 4, 6, 8, 10 or 12,
- 10 (k) nucleic acid sequences which are degenerated as a result of the genetic code to a nucleotide sequence of a nucleic acid as given in any of SEQ ID NOs 1, 3, 5, 7, 9 or 11, or as defined in (a) to (j),
- (l) nucleic acid sequences which are diverging due to the differences in codon usage between the organisms to a nucleotide sequence encoding a protein as given in any of SEQ ID NOs 2, 4, 6, 8, 10, 12 or as defined in 15 (a) to (j),
- (m) nucleic acid sequences which are diverging due to the differences between alleles encoding a protein as given in SEQ ID NOs 2, 4, 6, 8, 10 or 12, or as defined in (a) to (j),
- 20 (n) nucleic acid sequences encoding an immunologically active and/or functional fragment of a protein encoded by a DNA sequence as given in any of SEQ ID NOs 1, 3, 5, 7, 9 or 11, or a functional fragment of a nucleic acid as defined in any one of (a) to (m), and,
- 25 (o) nucleic acid sequences encoding a protein as defined in SEQ ID NO 2, 4, 6, 8, 10 or 12 or a nucleic acid as defined in any one of (a) to (n) characterized in that said sequence is a cDNA sequence.

The invention also relates to an isolated nucleic acid according to claim 2 which is DNA, cDNA, genomic DNA or synthetic DNA, or RNA wherein T is replaced by U.

The invention also relates to a nucleic acid molecule of at least 15 nucleotides in length hybridizing specifically with or specifically amplifying a nucleic acid of claim 2 or 3.

5 According to another embodiment, the invention also relates to a vector comprising a nucleic acid sequence according to claim 2 or 3. In a preferred embodiment, said vector is an expression vector wherein the nucleic acid sequence is operably linked to one or more control sequences allowing the expression of said sequence in prokaryotic and/or eukaryotic host cells.

10 It should be understood that for expression of the cytokinin oxidase genes of the invention in monocots, a nucleic acid sequence corresponding to the cDNA sequence should be used to avoid mis-splicing of introns in monocots.

The invention also relates to a host cell containing any of the nucleic acid molecules or vectors of the invention. Said host cell is chosen from the group comprising bacterial, insect, fungal, plant or animal cells.

15 Another embodiment of the invention relates to an isolated polypeptide encodable by a nucleic acid of claim 2 or 3, or a homologue or a derivative thereof, or an immunologically active and/or functional fragment thereof.

20 In a preferred embodiment, the invention relates to a polypeptide which has an amino acid sequence as given in SEQ ID NO 2, 4, 6, 8, 10 or 12, or a homologue or a derivative thereof, or an immunologically active and/or functional fragment thereof.

25 According to yet another embodiment, the invention relates to a method for producing a polypeptide according to claim 10 or 11 comprising culturing a host cell of claim 8 or 9 under conditions allowing the expression of the polypeptide and recovering the produced polypeptide from the culture.

The invention also relates to an antibody specifically recognizing a polypeptide of claim 10 or 11 or a specific epitope thereof.

30 The invention further relates to a method for the production of transgenic plants, plant cells or plant tissues comprising the introduction of a nucleic acid molecule according to any of claims 2 or 3 in an expressible format or a vector according to claim 6 or 7 in said plant, plant cell or plant tissue.

The invention also relates to a method for the production of altered plants, plant cells or plant tissues comprising the introduction of a polypeptide of claim 10 or 11 directly into a cell, a tissue or an organ of said plant.

5 According to another embodiment, the invention relates to a method for effecting the expression of a polypeptide of claim 10 or 11 comprising the introduction of a nucleic acid molecule of claim 2 or 3 operably linked to one or more control sequences or a vector according to claim 6 or 7 stably into the genome of a plant cell.

10 The invention further relates to a method of claim 15 or 16 further comprising regenerating a plant from said plant cell.

The invention also relates to a transgenic plant cell comprising a nucleic acid sequence of claim 2 or 3 which is operably linked to regulatory elements allowing transcription and/or expression of said nucleic acid in plant cells or obtainable by a method of claim 15 or 16.

15 According to a preferred embodiment, the invention relates to a transgenic plant cell of claim 18 wherein said nucleic acid of claim 2 or 3 is stably integrated into the genome of said plant cell.

The invention also relates to a transgenic plant or plant tissue comprising plant cells of claim 18 or 19 and to a harvestable part of said transgenic plant.

20 According to the invention, the harvestable part of said transgenic plant is selected from the group consisting of seeds, leaves, fruits, stem cultures, rhizomes and bulbs.

The invention also relates to the progeny derived from any of the plants or plant parts of any of claims 20 to 22.

25 According to another embodiment, the invention relates to a method for stimulating root growth comprising expression of a nucleic acid of claim 2 or 3 or comprising expression of another protein that reduces the level of active cytokinins in plants or plant parts.

30 The invention further relates to a method for enhancing the formation of lateral and adventitious roots comprising expression of a nucleic acid of claim 2 or 3 or

comprising expression of another protein that reduces the level of active cytokinins in plants or plant parts.

The invention also relates to method for altering root geotropism comprising altering the expression of a nucleic acid of claim 2 or 3 or comprising expression
5 of another protein that that reduces the level of active cytokinins in plants or plant parts.

The invention also relates to methods for enhancing early vigor and/or for modifying root/shoot ratio and/or for improving lodging and/or for increasing drought tolerance and/or for promoting *in vitro* propagation of explants
10 comprising expression of a nucleic acid of claim 2 or 3 or comprising expression of another protein that reduces the level of active cytokinins in plants or plant parts.

Furthermore, the invention relates to any of the above described methods, preferably a method of any of claims 24 to 26, said method leading to an
15 increase in yield.

The invention further relates to any of the above described methods, preferably a method of any of claims 24 to 27, wherein said expression of said nucleic acid occurs under the control of a strong constitutive promoter.

In a preferred embodiment the invention relates to any of the above described
20 methods, preferably a method of any of claims 24 to 27, wherein said expression of said nucleic acid occurs under the control of a promoter that is preferentially expressed in roots.

According to yet another embodiment, the invention relates to a method for modifying cell fate and/or modifying plant development and/or modifying plant
25 morphology and/or modifying plant biochemistry and/or modifying plant physiology and/or modifying the cell cycle progression rate comprising the modification of expression in particular cells, tissues or organs of a plant, of a nucleic acid sequence of claim 2 or 3.

The invention also relates to a method for obtaining enhanced growth, and/or
30 increased yield and/or delayed senescence of a plant cell, tissue and/or organ

and/or increased frequency of formation of lateral organs in a plant, comprising the ectopic expression of a nucleic acid sequence of claim 2 or 3.

The invention also relates to a method for promoting and extending cell division activity in cells in adverse growth conditions and/or in stress, comprising the
5 ectopic expression of a nucleic acid sequence of claim 2 or 3.

According to yet another embodiment, the invention relates to a method for identifying and obtaining proteins interacting with a polypeptide of claim 10 or 11 comprising a screening assay wherein a polypeptide of claim 10 or 11 is used.

In a more preferred embodiment, the invention relates to a method for identifying
10 and obtaining proteins interacting with a polypeptide of claim 10 or 11 comprising a two-hybrid screening assay wherein a polypeptide of claim 10 or 11 as a bait and a cDNA library as prey are used.

The invention further relates to a method for modulating the interaction between a polypeptide of claim 10 or 11 and interacting protein partners obtainable by a
15 method according to claim 30 or 31.

In a further embodiment, the invention relates to a method for identifying and obtaining compounds interacting with a polypeptide of claim 10 or 11 comprising the steps of:

- 20 a) providing a two-hybrid system wherein a polypeptide of claim 10 or 11 and an interacting protein partner obtainable by a method according to claim 30 or 31 are expressed,
- b) interacting said compound with the complex formed by the expressed polypeptides as defined in a), and,
- 25 c) performing (real-time) measurement of interaction of said compound with said polypeptide or the complex formed by the expressed polypeptides as defined in a).

The invention further relates to a method for identifying compounds or mixtures of compounds which specifically bind to a polypeptide of claim 10 or 11, comprising:

- 30 a) combining a polypeptide of claim 10 or 11 with said compound or mixtures of compounds under conditions suitable to allow complex formation, and,

- b) detecting complex formation, wherein the presence of a complex identifies a compound or mixture which specifically binds said polypeptide.

The invention also relates to a method of any of claims 30 to 34 wherein said compound or mixture inhibits the activity of said polypeptide of claim 10 or 11 and can be used for the rational design of chemicals.

According to another embodiment, the invention relates to the use of a compound or mixture identified by means of a method of any of claims 30 to 34 as a plant growth regulator or herbicide.

The invention also relates to a method for production of a plant growth regulator or herbicide composition comprising the steps of the method of any of claims 30 to 34 and formulating the compounds obtained from said steps in a suitable form for the application in agriculture or plant cell or tissue culture.

The invention also relates to a method for the design of or screening for growth-promoting chemicals or herbicides comprising the use of a nucleic acid of claim 2 or 3 or a vector of claim 6 or 7.

According to another embodiment, the invention relates to the use of a nucleic acid molecule of claim 2 or 3, the vector of claim 6 or 7, a polypeptide of claim 10 or 11 for increasing yield.

The invention also relates to the use of a nucleic acid molecule of claim 2 or 3, the vector of claim 6 or 7, a polypeptide of claim 10 or 11 for stimulating root growth.

The invention also relates to the use of a nucleic acid molecule of claim 2 or 3, the vector of claim 6 or 7, a polypeptide of claim 10 or 11 for enhancing the formation of lateral and adventitious roots.

The invention also relates to the use of a nucleic acid molecule of claim 2 or 3, the vector of claim 6 or 7, a polypeptide of claim 10 or 11 for altering root geotropism.

The invention further relates to the use of a nucleic acid molecule of claim 2 or 3, the vector of claim 6 or 7, a polypeptide of claim 10 or 11 for enhancing early vigor and/or for modifying root/shoot ratio and/or for improving lodging and/or for

increasing drought tolerance and/or for promoting *in vitro* propagation of explants.

The invention also relates to the use of a nucleic acid molecule of claim 2 or 3, the recombinant vector of claim 6 or 7, a polypeptide of claim 10 or 11 for
5 modifying plant development and/or for modifying plant morphology and/or for modifying plant biochemistry and/or for modifying plant physiology.

According to yet another embodiment, the invention relates to a diagnostic composition comprising at least a nucleic acid molecule of any of claims 2 to 5, the vector of claim 6 or 7, a polypeptide of claim 10 or 11 or an antibody of claim
10 13.

DEFINITIONS AND ELABORATIONS TO THE EMBODIMENTS

15 Those skilled in the art will be aware that the invention described herein is subject to variations and modifications other than those specifically described. It is to be understood that the invention described herein includes all such variations and modifications. The invention also includes all such steps, features, compositions and compounds referred to or indicated in this specification,
20 individually or collectively, and any and all combinations of any or more of said steps or features.

The present invention is applicable to any plant, in particular a monocotyledonous plants and dicotyledonous plants including a fodder or forage legume, ornamental plant, food crop, tree, or shrub selected from the list
25 comprising *Acacia spp.*, *Acer spp.*, *Actinidia spp.*, *Aesculus spp.*, *Agathis australis*, *Albizia amara*, *Alsophila tricolor*, *Andropogon spp.*, *Arachis spp.*, *Areca catechu*, *Astelia fragrans*, *Astragalus cicer*, *Baikiaea plurijuga*, *Betula spp.*, *Brassica spp.*, *Bruguiera gymnorhiza*, *Burkea africana*, *Butea frondosa*, *Cadaba farinosa*, *Calliandra spp.*, *Camellia sinensis*, *Canna indica*, *Capsicum spp.*,
30 *Cassia spp.*, *Centroema pubescens*, *Chaenomeles spp.*, *Cinnamomum cassia*, *Coffea arabica*, *Colophospermum mopane*, *Coronillia varia*, *Cotoneaster serotina*, *Crataegus spp.*, *Cucumis spp.*, *Cupressus spp.*, *Cyathea dealbata*,

- Cydonia oblonga*, *Cryptomeria japonica*, *Cymbopogon* spp., *Cynthea dealbata*,
Cydonia oblonga, *Dalbergia monetaria*, *Davallia divaricata*, *Desmodium* spp.,
Dicksonia squarosa, *Diheteropogon amplexans*, *Dioclea* spp., *Dolichos* spp.,
Dorycnium rectum, *Echinochloa pyramidalis*, *Ehrhartia* spp., *Eleusine coracana*,
5 *Eragrestis* spp., *Erythrina* spp., *Eucalyptus* spp., *Euclea schimperi*, *Eulalia*
villosa, *Fagopyrum* spp., *Feijoa sellowiana*, *Fragaria* spp., *Flemingia* spp.,
Freycinetia banksii, *Geranium thunbergii*, *Ginkgo biloba*, *Glycine javanica*,
Gliricidia spp., *Gossypium hirsutum*, *Grevillea* spp., *Guibourtia coleosperma*,
Hedysarum spp., *Hemarthia altissima*, *Heteropogon contortus*, *Hordeum vulgare*,
10 *Hyparrhenia rufa*, *Hypericum erectum*, *Hyperthelia dissoluta*, *Indigo incarnata*,
Iris spp., *Leptarrhena pyrolifolia*, *Lespedeza* spp., *Lettuca* spp., *Leucaena*
leucocephala, *Loudetia simplex*, *Lotonus bainesii*, *Lotus* spp., *Macrotyloma*
axillare, *Malus* spp., *Manihot esculenta*, *Medicago sativa*, *Metasequoia*
glyptostroboides, *Musa sapientum*, *Nicotianum* spp., *Onobrychis* spp.,
15 *Ornithopus* spp., *Oryza* spp., *Peltophorum africanum*, *Pennisetum* spp., *Persea*
gratissima, *Petunia* spp., *Phaseolus* spp., *Phoenix canariensis*, *Phormium*
cookianum, *Photinia* spp., *Picea glauca*, *Pinus* spp., *Pisum sativum*, *Podocarpus*
totara, *Pogonarthria fleckii*, *Pogonarthria squarrosa*, *Populus* spp., *Prosopis*
cineraria, *Pseudotsuga menziesii*, *Pterolobium stellatum*, *Pyrus communis*,
20 *Quercus* spp., *Rhaphiolepis umbellata*, *Rhopalostylis sapida*, *Rhus natalensis*,
Ribes grossularia, *Ribes* spp., *Robinia pseudoacacia*, *Rosa* spp., *Rubus* spp.,
Salix spp., *Schyzachyrium sanguineum*, *Sciadopitys verticillata*, *Sequoia*
sempervirens, *Sequoiadendron giganteum*, *Sorghum bicolor*, *Spinacia* spp.,
Sporobolus fimbriatus, *Stiburus alopecuroides*, *Stylosanthos humilis*, *Tadehagi*
25 *spp*, *Taxodium distichum*, *Themeda triandra*, *Trifolium* spp., *Triticum* spp., *Tsuga*
heterophylla, *Vaccinium* spp., *Vicia* spp., *Vitis vinifera*, *Watsonia pyramidata*,
Zantedeschia aethiopica, *Zea mays*, amaranth, artichoke, asparagus, broccoli,
brussel sprout, cabbage, canola, carrot, cauliflower, celery, collard greens, flax,
kale, lentil, oilseed rape, okra, onion, potato, rice, soybean, straw, sugarbeet,
30 sugar cane, sunflower, tomato, squash, and tea, amongst others, or the seeds of

any plant specifically named above or a tissue, cell or organ culture of any of the above species.

Throughout this specification, unless the context requires otherwise the word "comprise", and variations such as "comprises" and "comprising", will be understood to imply the inclusion of a stated integer or step or group of integers or steps but not the exclusion of any other integer or step or group of integers or steps.

As used herein, the term "derived from" shall be taken to indicate that a particular integer or group of integers has originated from the species specified, but has not necessarily been obtained directly from the specified source.

The terms "protein(s)", "peptide(s)" or "oligopeptide(s)", when used herein refer to amino acids in a polymeric form of any length. Said terms also include known amino acid modifications such as disulphide bond formation, cysteinylolation, oxidation, glutathionylation, methylation, acetylation, farnesylation, biotinylation, stearylation, formylation, lipoic acid addition, phosphorylation, sulphation, ubiquitination, myristoylation, palmitoylation, geranylgeranylation, cyclization (e.g. pyroglutamic acid formation), oxidation, deamidation, dehydration, glycosylation (e.g. pentoses, hexosamines, N-acetylhexosamines, deoxyhexoses, hexoses, sialic acid etc.) and acylation as well as non-naturally occurring amino acid residues, L-amino acid residues and D-amino acid residues.

"Homologues" of a protein of the invention are those peptides, oligopeptides, polypeptides, proteins and enzymes which contain amino acid substitutions, deletions and/or additions relative to the said protein with respect to which they are a homologue, without altering one or more of its functional properties, in particular without reducing the activity of the resulting. For example, a homologue of said protein will consist of a bioactive amino acid sequence variant of said protein. To produce such homologues, amino acids present in the said protein can be replaced by other amino acids having similar properties, for example hydrophobicity, hydrophilicity, hydrophobic moment, antigenicity, propensity to form or break α -helical structures or β -sheet structures, and so on. An overview of physical and chemical properties of amino acids is given in Table 1.

Table 1. Properties of naturally occurring amino acids.

Charge properties / hydrophobicity	Side group	Amino Acid
nonpolar hydrophobic	Aliphatic aliphatic, S-containing aromatic imino	ala, ile, leu, val met phe, trp pro
polar uncharged	Aliphatic amide aromatic hydroxyl sulfhydryl	gly asn, gln tyr ser, thr cys
positively charged	Basic	arg, his, lys
negatively charged	Acidic	asp, gly

5

Substitutional variants of a protein of the invention are those in which at least one residue in said protein amino acid sequence has been removed and a different residue inserted in its place. Amino acid substitutions are typically of single residues, but may be clustered depending upon functional constraints placed upon the polypeptide; insertions will usually be of the order of about 1-10 amino acid residues. and deletions will range from about 1-20 residues. Preferably, amino acid substitutions will comprise conservative amino acid substitutions, such as those described *supra*.

15 Insertional amino acid sequence variants of a protein of the invention are those in which one or more amino acid residues are introduced into a predetermined site in said protein. Insertions can comprise amino-terminal and/or carboxy-terminal fusions as well as intra-sequence insertions of single or multiple amino acids. Generally, insertions within the amino acid sequence will be smaller than amino or carboxyl terminal fusions, of the order of about 1 to 10 residues. Examples of

20

amino- or carboxy-terminal fusion proteins or peptides include the binding domain or activation domain of a transcriptional activator as used in a two-hybrid system, phage coat proteins, (histidine)₆-tag, glutathione S-transferase, protein A, maltose-binding protein, dihydrofolate reductase, Tag•100 epitope
5 (EETARFQPGYRS), c-myc epitope (EQKLISEEDL), FLAG[®]-epitope (DYKDDDK), lacZ, CMP (calmodulin-binding peptide), HA epitope (YPYDVPDYA), protein C epitope (EDQVDPRLIDGK) and VSV epitope (YTDIEMNRLGK).

Deletional variants of a protein of the invention are characterised by the removal
10 of one or more amino acids from the amino acid sequence of said protein.

Amino acid variants of a protein of the invention may readily be made using peptide synthetic techniques well known in the art, such as solid phase peptide synthesis and the like, or by recombinant DNA manipulations. The manipulation of DNA sequences to produce variant proteins which manifest as substitutional,
15 insertional or deletional variants are well known in the art. For example, techniques for making substitution mutations at predetermined sites in DNA having known sequence are well known to those skilled in the art, such as by M13 mutagenesis, T7-Gen in vitro mutagenesis kit (USB, Cleveland, OH), QuickChange Site Directed mutagenesis kit (Stratagene, San Diego, CA), PCR-
20 mediated site-directed mutagenesis or other site-directed mutagenesis protocols. In the current invention "homology" and "similarity" percentages between proteins are calculated using computer programs known in the art such as the Dnastar/MegAlign programs in combination with the Clustal method.

"Derivatives" of a protein of the invention are those peptides, oligopeptides,
25 polypeptides, proteins and enzymes which comprise at least about five contiguous amino acid residues of said polypeptide but which retain the biological activity of said protein. A "derivative" may further comprise additional naturally-occurring, altered glycosylated, acylated or non-naturally occurring amino acid residues compared to the amino acid sequence of a naturally-
30 occurring form of said polypeptide. Alternatively or in addition, a derivative may comprise one or more non-amino acid substituents compared to the amino acid

sequence of a naturally-occurring form of said polypeptide, for example a reporter molecule or other ligand, covalently or non-covalently bound to the amino acid sequence such as, for example, a reporter molecule which is bound thereto to facilitate its detection.

5 With "immunologically active" is meant that a molecule or specific fragments thereof such as specific epitopes or haptens are recognized by, i.e. bind to antibodies. Specific epitopes may be determined using, for example, peptide scanning techniques as described in Geysen *et al.* (1996) (Geysen, H.M., Rodda, S.J. and Mason, T.J. (1986). A priori delineation of a peptide which mimics a discontinuous antigenic determinant. *Mol. Immunol.* **23**, 709-715.).

10 The term "fragment of a sequence" or "part of a sequence" means a truncated sequence of the original sequence referred to. The truncated sequence (nucleic acid or protein sequence) can vary widely in length; the minimum size being a sequence of sufficient size to provide a sequence with at least a comparable function and/or activity or the original sequence referred to (e. g. "functional fragment"), while the maximum size is not critical. In some applications, the maximum size usually is not substantially greater than that required to provide the desired activity and/or function(s) of the original sequence. Typically, the truncated amino acid sequence will range from about 5 to about 60 amino acids

15 in length. More typically, however, the sequence will be a maximum of about 50 amino acids in length, preferably a maximum of about 60 amino acids. It is usually desirable to select sequences of at least about 10, 12 or 15 amino acids, up to a maximum of about 20 or 25 amino acids.

25 Functional fragments include those comprising an epitope which is specific for the proteins according to the invention. Preferred functional fragments have a length of at least, for example, 5, 10, 25, 100, 150 or 200 amino acids.

In the context of the current invention are embodied homologues, derivatives and/or immunologically active and/or functional fragments of inventive cytokinin oxidases as defined supra. Particularly preferred homologues, derivatives and/or

30 immunologically active and/or functional fragments of the cytokinin oxidase proteins which are contemplated for use in the current invention are derived from

plants, more specifically from *Arabidopsis thaliana*, even more specifically said cytokinin oxidases are the *Arabidopsis thaliana* (At)CKX, or are capable of being expressed therein. The present invention clearly contemplates the use of functional homologues or derivatives and/or immunologically active fragments of the AtCKX proteins and is not to be limited in application to the use of a
5 nucleotide sequence encoding one of said AtCKX proteins.

Any of said proteins, polypeptides, peptides and fragments thereof can be produced in a biological system, e.g. a cell culture. Alternatively any of said proteins, polypeptides, peptides and fragments thereof can be chemically
10 manufactured e.g. by solid phase peptide synthesis. Said proteins or fragments thereof can be part of a fusion protein as is the case in e.g. a two-hybrid assay which enables e.g. the identification of proteins interacting with a cytokinin oxidase according to the invention.

The proteins or fragments thereof are furthermore useful e.g. to modulate the
15 interaction between a cytokinin oxidase according to the invention and interacting protein partners obtained by a method of the invention. Chemically synthesized peptides are particularly useful e.g. as a source of antigens for the production of antisera and/or antibodies.

"Antibodies" include monoclonal, polyclonal, synthetic or heavy chain camel
20 antibodies as well as fragments of antibodies such as Fab, Fv or scFv fragments. Monoclonal antibodies can be prepared by the techniques as described in e.g. Liddle and Cryer (1991) which comprise the fusion of mouse myeloma cells to spleen cells derived from immunized animals. Furthermore, antibodies or fragments thereof to a molecule or fragments thereof can be obtained by using
25 methods as described in e.g. Harlow and Lane (1988). In the case of antibodies directed against small peptides such as fragments of a protein of the invention, said peptides are generally coupled to a carrier protein before immunization of animals. Such protein carriers include keyhole limpet hemocyanin (KLH), bovine serum albumin (BSA), ovalbumin and Tetanus toxoid. The carrier protein
30 enhances the immune response of the animal and provides epitopes for T-cell receptor binding sites. The term "antibodies" furthermore includes derivatives

thereof such as labelled antibodies. Antibody labels include alkaline phosphatase, PKH2, PKH26, PKH67, fluorescein (FITC), Hoechst 33258, R-phycoerythrin (PE), rhodamine (TRITC), Quantum Red, Texas Red, Cy3, biotin, agarose, peroxidase and gold spheres. Tools in molecular biology relying on antibodies against a protein include protein gel blot analysis, screening of expression libraries allowing gene identification, protein quantitative methods including ELISA and RIA, immunoaffinity purification of proteins, immunoprecipitation of proteins (see e.g. Example 5) and immunolocalization (see e.g. Example 6). Other uses of antibodies and especially of peptide antibodies include the study of proteolytic processing (Loffler et al. 1994, Woulfe et al. 1994), determination of protein active sites (Lerner 1982), the study of precursor and post-translational processing (Baron and Baltimore 1982, Lerner et al. 1981, Semier et al. 1982), identification of protein domains involved in protein-protein interactions (Murakami et al. 1992) and the study of exon usage in gene expression (Tamura et al. 1991).

Embodied in the current invention are antibodies specifically recognizing a cytokinin oxidase or homologue, derivative or fragment thereof as defined supra. Preferably said cytokinin oxidase is a plant cytokinin oxidase, more specifically one of the *Arabidopsis thaliana* cytokinin oxidases (AtCKX).

The terms "gene(s)", "polynucleotide(s)", "nucleic acid sequence(s)", "nucleotide sequence(s)", or "nucleic acid molecule(s)", when used herein refer to nucleotides, either ribonucleotides or deoxyribonucleotides or a combination of both, in a polymeric form of any length. Said terms furthermore include double-stranded and single-stranded DNA and RNA. Said terms also include known nucleotide modifications such as methylation, cyclization and 'caps' and substitution of one or more of the naturally occurring nucleotides with an analog such as inosine. Modifications of nucleotides include the addition of acridine, amine, biotin, cascade blue, cholesterol, Cy3®, Cy5®, Cy5.5® Dabcyl, digoxigenin, dinitrophenyl, Edans, 6-FAM, fluorescein, 3'-glyceryl, HEX, IRD-700, IRD-800, JOE, phosphate psoralen, rhodamine, ROX, thiol (SH), spacers, TAMRA, TET, AMCA-S®, SE, BODIPY®, Marina Blue®, Pacific Blue®, Oregon

Green[®], Rhodamine Green[®], Rhodamine Red[®], Rhodol Green[®] and Texas Red[®]. Polynucleotide backbone modifications include methylphosphonate, 2'-OMe-methylphosphonate RNA, phosphorothiorate, RNA, 2'-OMeRNA. Base modifications include 2-amino-dA, 2-aminopurine, 3'-(ddA), 3'dA(cordycepin), 7-deaza-dA, 8-Br-dA, 8-oxo-dA, N⁶-Me-dA, abasic site (dSpacer), biotin dT, 2'-OMe-5Me-C, 2'-OMe-propynyl-C, 3'-(5-Me-dC), 3'-(ddC), 5-Br-dC, 5-I-dC, 5-Me-dC, 5-F-dC, carboxy-dT, convertible dA, convertible dC, convertible dG, convertible dT, convertible dU, 7-deaza-dG, 8-Br-dG, 8-oxo-dG, O⁶-Me-dG, S6-DNP-dG, 4-methyl-indole, 5-nitroindole, 2'-OMe-inosine, 2'-dI, O⁶-phenyl-dI, 4-methyl-indole, 2'-deoxynebularine, 5-nitroindole, 2-aminopurine, dP(purine analogue), dK(pyrimidine analogue), 3-nitropyrrole, 2-thio-dT, 4-thio-dT, biotin-dT, carboxy-dT, O⁴-Me-dT, O⁴-triazol dT, 2'-OMe-propynyl-U, 5-Br-dU, 2'-dU, 5-F-dU, 5-I-dU, O⁴-triazol dU. Said terms also encompass peptide nucleic acids (PNAs), a DNA analogue in which the backbone is a pseudopeptide consisting of N-(2-aminoethyl)-glycine units rather than a sugar. PNAs mimic the behaviour of DNA and bind complementary nucleic acid strands. The neutral backbone of PNA results in stronger binding and greater specificity than normally achieved. In addition, the unique chemical, physical and biological properties of PNA have been exploited to produce powerful biomolecular tools, antisense and antigene agents, molecular probes and biosensors.

The present invention also advantageously provides nucleic acid sequences of at least approximately 15 contiguous nucleotides of a nucleic acid according to the invention and preferably from 15 to 50 nucleotides. These sequences may, advantageously be used as probes to specifically hybridise to sequences of the invention as defined above or primers to initiate specific amplification or replication of sequences of the invention as defined above, or the like. Such nucleic acid sequences may be produced according to techniques well known in the art, such as by recombinant or synthetic means. They may also be used in diagnostic kits or the like for detecting the presence of a nucleic acid according to the invention. These tests generally comprise contacting the probe with the

sample under hybridising conditions and detecting the presence of any duplex or triplex formation between the probe and any nucleic acid in the sample.

Advantageously, the nucleic acid sequences, according to the invention may be produced using such recombinant or synthetic means, such as for example using

5 PCR cloning mechanisms which generally involve making a pair of primers, which may be from approximately 15 to 50 nucleotides to a region of the gene which is desired to be cloned, bringing the primers into contact with mRNA, cDNA or genomic DNA from a human cell, performing a polymerase chain reaction under conditions which bring about amplification of the desired region,
10 isolated the amplified region or fragment and recovering the amplified DNA. Generally, such techniques as defined herein are well known in the art, such as described in Sambrook et al. (Molecular Cloning: a Laboratory Manual, 1989).

A "coding sequence" or "open reading frame" or "ORF" is defined as a nucleotide sequence that can be transcribed into mRNA and/or translated into a polypeptide
15 when placed under the control of appropriate control sequences or regulatory sequences, i.e. when said coding sequence or ORF is present in an expressible format. Said coding sequence or ORF is bounded by a 5' translation start codon and a 3' translation stop codon. A coding sequence or ORF can include, but is not limited to RNA, mRNA, cDNA, recombinant nucleotide sequences,
20 synthetically manufactured nucleotide sequences or genomic DNA. Said coding sequence or ORF can be interrupted by intervening nucleic acid sequences.

Genes and coding sequences essentially encoding the same protein but isolated from different sources can consist of substantially divergent nucleic acid sequences. Reciprocally, substantially divergent nucleic acid sequences can be
25 designed to effect expression of essentially the same protein. Said nucleic acid sequences are the result of e.g. the existence of different alleles of a given gene, of the degeneracy of the genetic code or of differences in codon usage. Thus, as indicated in Table 2, amino acids such as methionine and tryptophan are encoded by a single codon whereas other amino acids such as arginine, leucine
30 and serine can each be translated from up to six different codons. Differences in preferred codon usage are illustrated in Table 3 for *Agrobacterium tumefaciens*

(a bacterium), *A. thaliana*, *M. sativa* (two dicotyledonous plants) and *Oryza sativa* (a monocotyledonous plant). To extract one example, the codon GGC (for glycine) is the most frequently used codon in *A. tumefaciens* (36.2 ‰), is the second most frequently used codon in *O. sativa* but is used at much lower frequencies in *A. thaliana* and *M. sativa* (9 ‰ and 8.4 ‰, respectively). Of the four possible codons encoding glycine (see Table 2), said GGC codon is most preferably used in *A. tumefaciens* and *O. sativa*. However, in *A. thaliana* this is the GGA (and GGU) codon whereas in *M. sativa* this is the GGU (and GGA) codon.

"Hybridization" is the process wherein substantially homologous complementary nucleotide sequences anneal to each other. The hybridization process can occur entirely in solution, i.e. both complementary nucleic acids are in solution. Tools in molecular biology relying on such a process include PCR, subtractive hybridization and DNA sequence determination. The hybridization process can also occur with one of the complementary nucleic acids immobilized to a matrix such as magnetic beads, Sepharose beads or any other resin. Tools in molecular biology relying on such a process include the isolation of poly (A+)mRNA. The hybridization process can furthermore occur with one of the complementary nucleic acids immobilized to a solid support such as a nitrocellulose or nylon membrane or immobilized by e.g. photolithography to e.g. a silicious glass support (the latter known as nucleic acid arrays or microarrays or as nucleic acid chips). Tools in molecular biology relying on such a process include RNA and DNA gel blot analysis, colony hybridization, plaque hybridization and microarray hybridization. In order to allow hybridization to occur, the nucleic acid molecules are generally thermally or chemically (e.g. by NaOH) denatured to melt a double strand into two single strands and/or to remove hairpins or other secondary structures from single stranded nucleic acids. The stringency of hybridization is influenced by conditions such as temperature, salt concentration and hybridization buffer composition. High stringency conditions for hybridization include high temperature and/or low salt concentration (salts include NaCl and Na₃-citrate) and/or the inclusion of formamide in the hybridization buffer and/or

lowering the concentration of compounds such as SDS (detergent) in the hybridization buffer and/or exclusion of compounds such as dextran sulfate or polyethylene glycol (promoting molecular crowding) from the hybridization buffer. Conventional hybridization conditions are described in e.g. Sambrook et al.
5 (1989) but the skilled craftsman will appreciate that numerous different hybridization conditions can be designed in function of the known or the expected homology and/or length of the nucleic acid sequence. Sufficiently low stringency hybridization conditions are particularly preferred to isolate nucleic acids heterologous to the DNA sequences of the invention defined supra. Elements
10 contributing to said heterology include allelism, degeneration of the genetic code and differences in preferred codon usage as discussed supra.

Clearly, the current invention embodies the use of the inventive DNA sequences encoding a cytokinin oxidase, homologue, derivative or immunologically active and/or functional fragment thereof as defined higher in any method of
15 hybridization . The current invention furthermore also related to DNA sequences hybridizing to said inventive DNA sequences. Preferably said cytokinin oxidase is a plant cytokinin oxidase, more specifically the *Arabidopsis thaliana* (At)CKX.

DNA sequences as defined in the current invention can be interrupted by intervening sequences. With "intervening sequences" is meant any nucleic acid
20 sequence which disrupts a coding sequence comprising said inventive DNA sequence or which disrupts the expressible format of a DNA sequence comprising said inventive DNA sequence. Removal of the intervening sequence restores said coding sequence or said expressible format. Examples of intervening sequences include introns and mobilizable DNA sequences such as
25 transposons. With "mobilizable DNA sequence" is meant any DNA sequence that can be mobilized as the result of a recombination event.

Table 2. Degeneracy of the genetic code.

Amino Acid	Three-letter code	One-letter code	Possible codons					
Alanine	Ala	A	GCA	GCC	GCG	GCU		
Arginine	Arg	R	AGA	AGG	CGA	CGC	CGG	CGU
Asparagine	Asn	N	AAC	AAU				
Aspartic Acid	Asp	D	GAC	GAU				
Cysteine	Cys	C	UGC	UGU				
Glutamic Acid	Glu	E	GAA	GAG				
Glutamine	Gln	Q	CAA	CAG				
Glycine	Gly	G	GGA	GGC	GGG	GGU		
Histidine	His	H	CAC	CAU				
Isoleucine	Ile	I	AUA	AUC	AUU			
Leucine	Leu	L	UUA	UUG	CUA	CUC	CUG	CUU
Lysine	Lys	K	AAA	AAG				
Methionine	Met	M	AUG					
Phenylalanine	Phe	F	UUC	UUU				
Proline	Pro	P	CCA	CCC	CCG	CCU		
Serine	Ser	S	AGC	AGU	UCA	UCC	UCG	UCU
Threonine	Thr	T	ACA	ACC	ACG	ACU		
Tryptophan	Trp	W	UGG					
Tyrosine	Tyr	Y	UAC	UAU				
Valine	Val	V	GUA	GUC	GUG	GUU		
Possible "STOP" codons								
UAA	UAG	UGA						

5

Table 3. Usage of the indicated codons in the different organisms given as frequency per thousand codons (<http://www.kazusa.or.jp/codon>).

Codon	<i>Agrobacterium tumefaciens</i>	<i>Arabidopsis thaliana</i>	<i>Medicago sativa</i>	<i>Oryza sativa</i>
UUU	13.9	22.5	24.1	11.3
UUC	24.3	20.7	16.9	26.3
UUA	3.5	12.9	10.4	4.7
UUG	13.2	21.0	22.4	11.8
UCU	7.0	24.6	19.8	10.1
UCC	14.8	10.8	7.7	16.9
UCA	7.4	17.8	17.2	9.7
UCG	18.2	8.9	3.2	10.8
UAU	12.3	15.2	16.6	9.2
UAC	10.3	13.7	14.0	20.6
UAA	0.9	0.9	1.2	0.9

UAG	0.6	0.5	0.8	0.8
UGU	3.0	10.8	10.6	5.0
UGC	7.4	7.2	5.8	14.3
UGA	1.8	1.0	0.8	1.3
UGG	12.2	12.7	10.0	12.8
CUU	19.1	24.3	28.3	14.6
CUC	25.7	15.9	12.0	28.0
CUA	5.2	10.0	8.8	5.7
CUG	31.6	9.9	8.5	22.1
CCU	7.7	18.3	23.2	11.8
CCC	10.6	5.3	5.3	12.5
CCA	8.9	16.1	22.6	12.2
CCG	20.7	8.3	3.6	16.7
CAU	10.6	14.0	14.6	9.2
CAC	9.1	8.7	9.1	14.6
CAA	11.2	19.7	23.2	11.9
CAG	24.9	15.2	12.3	24.6
CGU	12.2	8.9	10.1	6.8
CGC	25.5	3.7	4.2	15.9
CGA	8.2	6.2	4.2	4.2
CGG	13.2	4.8	1.8	9.7
AUU	15.4	22.0	29.4	13.8
AUC	36.9	18.5	14.7	25.5
AUA	6.2	12.9	11.7	7.2
AUG	24.7	24.5	21.7	24.4
ACU	6.4	17.8	20.8	10.3
ACC	20.9	10.3	11.7	18.6
ACA	9.1	15.9	18.9	10.0
ACG	18.8	7.6	2.8	10.8
AAU	13.5	22.7	25.0	12.9
AAC	18.7	20.9	18.7	25.1
AAA	13.6	31.0	32.2	12.0
AAG	24.4	32.6	35.1	39.4
AGU	5.7	14.0	12.6	7.3
AGC	15.8	11.1	8.8	16.9
AGA	5.3	18.7	13.6	7.7
AGG	6.5	10.9	11.7	14.9
GUU	16.6	27.3	34.7	15.0
GUC	29.3	12.7	9.9	22.8
GUA	6.1	10.1	10.0	5.7
GUG	19.7	17.5	16.5	25.0
GCU	17.4	28.0	34.6	19.8
GCC	35.8	10.3	11.4	33.2
GCA	19.5	17.6	25.9	15.6
GCG	31.7	8.8	3.4	25.3
GAU	25.8	36.8	40.0	21.5
GAC	28.0	17.3	15.5	31.6
GAA	29.9	34.4	35.9	17.1
GAG	26.3	32.2	27.4	41.1

GGU	16.5	22.2	28.7	16.3
GGC	36.2	9.0	8.4	34.7
GGA	12.5	23.9	27.3	15.0
GGG	11.3	10.2	7.4	16.6

To effect expression of a protein in a cell, tissue or organ, preferably of plant origin, either the protein may be introduced directly to said cell, such as by microinjection or ballistic means or alternatively, an isolated nucleic acid molecule encoding said protein may be introduced into said cell, tissue or organ in an expressible format.

Preferably, the DNA sequence of the invention comprises a coding sequence or open reading frame (ORF) encoding a cytokinin oxidase protein or a homologue or derivative thereof or an immunologically active and/or functional fragment thereof as defined supra. The preferred protein of the invention comprises the amino acid sequence of said cytokinin oxidase. Preferably said cytokinin oxidase is a plant cytokinin oxidase and more specifically a *Arabidopsis thaliana* (At)CKX. With "vector" or "vector sequence" is meant a DNA sequence which can be introduced in an organism by transformation and can be stably maintained in said organism. Vector maintenance is possible in e.g. cultures of *Escherichia coli*, *A. tumefaciens*, *Saccharomyces cerevisiae* or *Schizosaccharomyces pombe*. Other vectors such as phagemids and cosmid vectors can be maintained and multiplied in bacteria and/or viruses. Vector sequences generally comprise a set of unique sites recognized by restriction enzymes, the multiple cloning site (MCS), wherein one or more non-vector sequence(s) can be inserted.

With "non-vector sequence" is accordingly meant a DNA sequence which is integrated in one or more of the sites of the MCS comprised within a vector.

"Expression vectors" form a subset of vectors which, by virtue of comprising the appropriate regulatory or control sequences enabling the creation of an expressible format for the inserted non-vector sequence(s), thus allowing expression of the protein encoded by said non-vector sequence(s). Expression vectors are known in the art enabling protein expression in organisms including bacteria (e.g. *E. coli*), fungi (e.g. *S. cerevisiae*, *S. pombe*, *Pichia pastoris*), insect

cells (e.g. baculoviral expression vectors), animal cells (e.g. COS or CHO cells) and plant cells (e.g. potato virus X-based expression vectors).

The current invention clearly includes any cytokinin oxidase, homologue, derivative and/or immunologically active and/or functional fragment thereof as defined supra. Preferably said cytokinin oxidase is a plant cytokinin oxidase, more specifically a *Arabidopsis thaliana* (At)CKX.

As an alternative to expression vector-mediated protein production in biological systems, chemical protein synthesis can be applied. Synthetic peptides can be manufactured in solution phase or in solid phase. Solid phase peptide synthesis (Merrifield 1963) is, however, the most common way and involves the sequential addition of amino acids to create a linear peptide chain. Solid phase peptide synthesis includes cycles consisting of three steps: (i) immobilization of the carboxy-terminal amino acid of the growing peptide chain to a solid support or resin; (ii) chain assembly, a process consisting of activation, coupling and deprotection of the amino acid to be added to the growing peptide chain; and (iii) cleavage involving removal of the completed peptide chain from the resin and removal of the protecting groups from the amino acid side chains. Common approaches in solid phase peptide synthesis include Fmoc/tBu (9-fluorenylmethyloxycarbonyl/t-butyl) and Boc (t-butyloxycarbonyl) as the amino-terminal protecting groups of amino acids. Amino acid side chain protecting groups include methyl (Me), formyl (CHO), ethyl (Et), acetyl (Ac), t-butyl (t-Bu), anisyl, benzyl (Bzl), trifluoroacetyl (Tfa), N-hydroxysuccinimide (ONSu, OSu), benzoyl (Bz), 4-methylbenzyl (Meb), thioanisyl, thiocresyl, benzyloxymethyl (Bom), 4-nitrophenyl (ONp), benzyloxycarbonyl (Z), 2-nitrobenzoyl (NBz), 2-nitrophenylsulphenyl (Nps), 4-toluenesulphonyl (Tosyl, Tos), pentafluorophenyl (Pfp), diphenylmethyl (Dpm), 2-chlorobenzyloxycarbonyl (Cl-Z), 2,4,5-trichlorophenyl, 2-bromobenzyloxycarbonyl (Br-Z), triphenylmethyl (Trityl, Trt), and 2,5,7,8-pentamethyl-chroman-6-sulphonyl (Pmc). During chain assembly, Fmoc or Boc are removed resulting in an activated amino-terminus of the amino acid residue bound to the growing chain. The carboxy-terminus of the incoming amino acid is activated by conversion into a highly reactive ester, e.g. by HBTU. With

current technologies (e.g. PerSeptive Biosystems 9050 synthesizer, Applied Biosystems Model 431A Peptide Synthesizer), linear peptides of up to 50 residues can be manufactured. A number of guidelines is available to produce peptides that are suitable for use in biological systems including (i) limiting the
5 use of difficult amino acids such as cys, met, trp (easily oxidized and/or degraded during peptide synthesis) or arg; (ii) minimize hydrophobic amino acids (can impair peptide solubility); and (iii) prevent an amino-terminal glutamic acid (can cyclize to pyroglutamate).

By "expressible format" is meant that the isolated nucleic acid molecule is in a
10 form suitable for being transcribed into mRNA and/or translated to produce a protein, either constitutively or following induction by an intracellular or extracellular signal, such as an environmental stimulus or stress (mitogens, anoxia, hypoxia, temperature, salt, light, dehydration, etc) or a chemical compound such as IPTG (isopropyl- β -D-thiogalactopyranoside) or such as an
15 antibiotic (tetracycline, ampicillin, rifampicin, kanamycin), hormone (e.g. gibberellin, auxin, cytokinin, glucocorticoid, brassinosteroid, ethylene, abscisic acid etc), hormone analogue (iodoacetic acid (IAA), 2,4-D, etc) , metal (zinc, copper, iron, etc), or dexamethasone, amongst others. As will be known to those skilled in the art, expression of a functional protein may also require one or more
20 post-translational modifications, such as glycosylation, phosphorylation, dephosphorylation, or one or more protein-protein interactions, amongst others. All such processes are included within the scope of the term "expressible format".

Preferably, expression of a protein in a specific cell, tissue, or organ, preferably
25 of plant origin, is effected by introducing and expressing an isolated nucleic acid molecule encoding said protein, such as a cDNA molecule, genomic gene, synthetic oligonucleotide molecule, mRNA molecule or open reading frame, to said cell, tissue or organ, wherein said nucleic acid molecule is placed operably in connection with suitable regulatory or control sequences including a promoter,
30 preferably a plant-expressible promoter, and a terminator sequence.

Reference herein to a "promoter" is to be taken in its broadest context and includes the transcriptional regulatory sequences derived from a classical eukaryotic genomic gene, including the TATA box which is required for accurate transcription initiation, with or without a CCAAT box sequence and additional
5 regulatory or control elements (i.e. upstream activating sequences, enhancers and silencers) which alter gene expression in response to developmental and/or external stimuli, or in a tissue-specific manner.

The term "promoter" also includes the transcriptional regulatory sequences of a classical prokaryotic gene, in which case it may include a -35 box sequence
10 and/or a -10 box transcriptional regulatory sequences.

The term "promoter" is also used to describe a synthetic or fusion molecule, or derivative which confers, activates or enhances expression of a nucleic acid molecule in a cell, tissue or organ.

Promoters may contain additional copies of one or more specific regulatory
15 elements, to further enhance expression and/or to alter the spatial expression and/or temporal expression of a nucleic acid molecule to which it is operably connected. Such regulatory elements may be placed adjacent to a heterologous promoter sequence to drive expression of a nucleic acid molecule in response to e.g. copper, glucocorticoids, dexamethasone, tetracycline, gibberellin, cAMP,
20 abscisic acid, auxin, wounding, ethylene, jasmonate or salicylic acid or to confer expression of a nucleic acid molecule to specific cells, tissues or organs such as meristems, leaves, roots, embryo, flowers, seeds or fruits.

In the context of the present invention, the promoter preferably is a plant-expressible promoter sequence. Promoters, however, that also function or solely
25 function in non-plant cells such as bacteria, yeast cells, insect cells and animal cells are not excluded from the invention. By "plant-expressible" is meant that the promoter sequence, including any additional regulatory elements added thereto or contained therein, is at least capable of inducing, conferring, activating or enhancing expression in a plant cell, tissue or organ, preferably a
30 monocotyledonous or dicotyledonous plant cell, tissue, or organ.

Regulatable promoters as part of a binary viral plant expression system are also known to the skilled artisan (Yadav 1999 – WO9922003; Yadav 2000 – WO0017365).

5 The terms "plant-operable" and "operable in a plant" when used herein, in respect of a promoter sequence, shall be taken to be equivalent to a plant-expressible promoter sequence.

10 In the present context, a "regulatable promoter sequence" is a promoter that is capable of conferring expression on a structural gene in a particular cell, tissue, or organ or group of cells, tissues or organs of a plant, optionally under specific conditions, however does generally not confer expression throughout the plant under all conditions. Accordingly, a regulatable promoter sequence may be a promoter sequence that confers expression on a gene to which it is operably connected in a particular location within the plant or alternatively, throughout the plant under a specific set of conditions, such as following induction of gene
15 expression by a chemical compound or other elicitor.

20 Preferably, the regulatable promoter used in the performance of the present invention confers expression in a specific location within the plant, either constitutively or following induction, however not in the whole plant under any circumstances. Included within the scope of such promoters are cell-specific promoter sequences, tissue-specific promoter sequences, organ-specific promoter sequences, cell cycle specific gene promoter sequences, inducible promoter sequences and constitutive promoter sequences that have been modified to confer expression in a particular part of the plant at any one time, such as by integration of said constitutive promoter within a transposable genetic
25 element (*Ac*, *Ds*, *Spm*, *En*, or other transposon).

The term "cell-specific" shall be taken to indicate that expression is predominantly in a particular cell or cell-type, preferably of plant origin, albeit not necessarily exclusively in said cell or cell-type.

30 Similarly, the term "tissue-specific" shall be taken to indicate that expression is predominantly in a particular tissue or tissue-type, preferably of plant origin, albeit not necessarily exclusively in said tissue or tissue-type.

Similarly, the term "organ-specific" shall be taken to indicate that expression is predominantly in a particular organ, preferably of plant origin, albeit not necessarily exclusively in said organ.

5 Similarly, the term "cell cycle specific" shall be taken to indicate that expression is predominantly cyclic and occurring in one or more, not necessarily consecutive phases of the cell cycle albeit not necessarily exclusively in cycling cells, preferably of plant origin.

Those skilled in the art will be aware that an "inducible promoter" is a promoter the transcriptional activity of which is increased or induced in response to a
10 developmental, chemical, environmental, or physical stimulus. Similarly, the skilled craftsman will understand that a "constitutive promoter" is a promoter that is transcriptionally active throughout most, but not necessarily all parts of an organism, preferably a plant, during most, but not necessarily all phases of its growth and development.

15 Those skilled in the art will readily be capable of selecting appropriate promoter sequences for use in regulating appropriate expression of the cytokinin oxidase protein from publicly-available or readily-available sources, without undue experimentation.

Placing a nucleic acid molecule under the regulatory control of a promoter
20 sequence, or in operable connection with a promoter sequence, means positioning said nucleic acid molecule such that expression is controlled by the promoter sequence. A promoter is usually, but not necessarily, positioned upstream, or at the 5'-end, and within 2 kb of the start site of transcription, of the nucleic acid molecule which it regulates. In the construction of heterologous
25 promoter/structural gene combinations it is generally preferred to position the promoter at a distance from the gene transcription start site that is approximately the same as the distance between that promoter and the gene it controls in its natural setting (i.e., the gene from which the promoter is derived). As is known in the art, some variation in this distance can be accommodated without loss of
30 promoter function. Similarly, the preferred positioning of a regulatory sequence element with respect to a heterologous gene to be placed under its control is

defined by the positioning of the element in its natural setting (i.e., the gene from which it is derived). Again, as is known in the art, some variation in this distance can also occur.

5 Examples of promoters suitable for use in gene constructs of the present invention include those listed in Table 4, amongst others. The promoters listed in Table 4 are provided for the purposes of exemplification only and the present invention is not to be limited by the list provided therein. Those skilled in the art will readily be in a position to provide additional promoters that are useful in performing the present invention.

10 In the case of constitutive promoters or promoters that induce expression throughout the entire plant, it is preferred that such sequences are modified by the addition of nucleotide sequences derived from one or more of the tissue-specific promoters listed in Table 4, or alternatively, nucleotide sequences derived from one or more of the above-mentioned tissue-specific inducible
15 promoters, to confer tissue-specificity thereon. For example, the CaMV 35S promoter may be modified by the addition of maize *Adh1* promoter sequence, to confer anaerobically-regulated root-specific expression thereon, as described previously (Ellis *et al.*, 1987). Another example describes conferring root specific or root abundant gene expression by fusing the CaMV35S promoter to elements
20 of the maize glycine-rich protein GRP3 gene (Feix and Wulff 2000 - WO0015662). Such modifications can be achieved by routine experimentation by those skilled in the art.

The term "terminator" refers to a DNA sequence at the end of a transcriptional unit which signals termination of transcription. Terminators are 3'-non-translated
25 DNA sequences containing a polyadenylation signal, which facilitates the addition of polyadenylate sequences to the 3'-end of a primary transcript. Terminators active in cells derived from viruses, yeasts, moulds, bacteria, insects, birds, mammals and plants are known and described in the literature. They may be isolated from bacteria, fungi, viruses, animals and/or plants.

Table 4. Exemplary plant-expressible promoters for use in the performance of the present invention

I: CELL-SPECIFIC, TISSUE-SPECIFIC, AND ORGAN-SPECIFIC PROMOTERS		
GENE SOURCE	EXPRESSION PATTERN	REFERENCE
α -amylase (<i>Amy32b</i>)	aleurone	Lanahan, M.B., <i>et al.</i> , <i>Plant Cell</i> 4:203-211, 1992; Skriver, K., <i>et al. Proc. Natl. Acad. Sci. (USA)</i> 88: 7266-7270, 1991
cathepsin β -like gene	aleurone	Cejudo, F.J., <i>et al. Plant Molecular Biology</i> 20:849-856, 1992.
<i>Agrobacterium rhizogenes rolB</i>	cambium	Nilsson <i>et al.</i> , <i>Physiol. Plant.</i> 100:456-462, 1997
AtPRP4	flowers	http://salus.medium.edu/mmg/tierney/html
chalcone synthase (<i>chsA</i>)	flowers	Van der Meer, <i>et al.</i> , <i>Plant Mol. Biol.</i> 15, 95-109, 1990.
LAT52	anther	Twell <i>et al</i> <i>Mol. Gen Genet.</i> 217:240-245 (1989)
<i>apetala-3</i>	flowers	
chitinase	fruit (berries, grapes, etc)	Thomas <i>et al.</i> CSIRO Plant Industry, Urrbrae, South Australia, Australia; http://winetitles.com.au/gwrdc/csh95-1.html
<i>rbcS-3A</i>	green tissue (eg leaf)	Lam, E. <i>et al.</i> , <i>The Plant Cell</i> 2: 857-866, 1990.; Tucker <i>et al.</i> , <i>Plant Physiol.</i> 113: 1303-1308, 1992.
leaf-specific genes	leaf	Baszczynski, <i>et al.</i> , <i>Nucl. Acid Res.</i> 16: 4732, 1988.
AtPRP4	leaf	http://salus.medium.edu/mmg/tierney/html
chlorella virus adenine methyltransferase gene promoter	leaf	Mitra and Higgins, 1994, <i>Plant Molecular Biology</i> 26: 85-93
aldP gene promoter from rice	leaf	Kagaya <i>et al.</i> , 1995, <i>Molecular and General Genetics</i> 248: 668-674
<i>rbcS</i> promoter from rice or tomato	leaf	Kyozuka <i>et al.</i> , 1993, <i>Plant Physiology</i> 102: 991-1000
<i>Pinus cab-6</i>	leaf	Yamamoto <i>et al.</i> , <i>Plant Cell Physiol.</i> 35:773-778, 1994.
rubisco promoter	leaf	

cab (chlorophyll a/b-binding protein)	leaf	
SAM22	senescent leaf	Crowell, et al., <i>Plant Mol. Biol.</i> 18: 459-466, 1992.
<i>ltp</i> gene (lipid transfer gene)		Fleming, et al, <i>Plant J.</i> 2, 855-862.
<i>R. japonicum nif</i> gene	Nodule	United States Patent No. 4, 803, 165
<i>B. japonicum nifH</i> gene	Nodule	United States Patent No. 5, 008, 194
GmENOD40	Nodule	Yang, et al., <i>The Plant J.</i> 3: 573-585.
PEP carboxylase (PEPC)	Nodule	Pathirana, et al., <i>Plant Mol. Biol.</i> 20: 437-450, 1992.
leghaemoglobin (Lb)	Nodule	Gordon, et al., <i>J. Exp. Bot.</i> 44: 1453-1465, 1993.
<i>Tungro bacilliform</i> virus gene	phloem	Bhattacharyya-Pakrasi, et al, <i>The Plant J.</i> 4: 71-79, 1992.
pollen-specific genes	pollen; microspore	Albani, et al., <i>Plant Mol. Biol.</i> 15: 605, 1990; Albani, et al., <i>Plant Mol. Biol.</i> 16: 501, 1991)
Zm13	pollen	Guerrero et al <i>Mol. Gen. Genet.</i> 224:161-168 (1993)
apg gene	microspore	Twell et al <i>Sex. Plant Reprod.</i> 6:217-224 (1993)
maize pollen-specific gene	pollen	Hamilton, et al., <i>Plant Mol. Biol.</i> 18: 211-218, 1992.
sunflower pollen-expressed gene	pollen	Baltz, et al., <i>The Plant J.</i> 2: 713-721, 1992.
<i>B. napus</i> pollen-specific gene	pollen;anther; tapetum	Arnoldo, et al., <i>J. Cell. Biochem.</i> , Abstract No. Y101, 204, 1992.
root-expressible genes	roots	Tingey, et al., <i>EMBO J.</i> 6: 1, 1987.
tobacco auxin-inducible gene	root tip	Van der Zaal, et al., <i>Plant Mol. Biol.</i> 16, 983, 1991.
β -tubulin	root	Oppenheimer, et al., <i>Gene</i> 63: 87, 1988.
tobacco root-specific genes	root	Conkling, et al., <i>Plant Physiol.</i> 93: 1203, 1990.
<i>B. napus</i> G1-3b gene	root	United States Patent No. 5, 401, 836
SbPRP1	roots	Suzuki et al., <i>Plant Mol. Biol.</i> 21: 109-119, 1993.
AtPRP1; AtPRP3	roots; root hairs	http://salus.medium.edu/mmg/tierney/html
RD2 gene	root cortex	http://www2.cnsu.edu/ncsu/research
TobRB7 gene	root vasculature	http://www2.cnsu.edu/ncsu/research
AtPRP4	leaves; flowers; lateral root primordia	http://salus.medium.edu/mmg/tierney/html

seed-specific genes	seed	Simon, <i>et al.</i> , <i>Plant Mol. Biol.</i> 5: 191, 1985; Scofield, <i>et al.</i> , <i>J. Biol. Chem.</i> 262: 12202, 1987.; Baszczynski, <i>et al.</i> , <i>Plant Mol. Biol.</i> 14: 633, 1990.
Brazil Nut albumin	seed	Pearson, <i>et al.</i> , <i>Plant Mol. Biol.</i> 18: 235-245, 1992.
legumin	seed	Ellis, <i>et al.</i> , <i>Plant Mol. Biol.</i> 10: 203-214, 1988.
glutelin (rice)	seed	Takaiwa, <i>et al.</i> , <i>Mol. Gen. Genet.</i> 208: 15-22, 1986; Takaiwa, <i>et al.</i> , <i>FEBS Letts.</i> 221: 43-47, 1987.
zein	seed	Matzke <i>et al</i> <i>Plant Mol Biol</i> , 14(3):323-32 1990
napA	seed	Stalberg, <i>et al</i> , <i>Planta</i> 199: 515-519, 1996.
wheat LMW and HMW glutenin-1	endosperm	<i>Mol Gen Genet</i> 216:81-90, 1989; <i>NAR</i> 17:461-2, 1989
wheat SPA	seed	Albani <i>et al</i> , <i>Plant Cell</i> , 9: 171-184, 1997
wheat α , β , γ -gliadins	endosperm	<i>EMBO</i> 3:1409-15, 1984
barley <i>ltr1</i> promoter	endosperm	
barley B1, C, D, hordein	endosperm	<i>Theor Appl Gen</i> 98:1253-62, 1999; <i>Plant J</i> 4:343-55, 1993; <i>Mol Gen Genet</i> 250:750-60, 1996
barley DOF	endosperm	Mena <i>et al</i> , <i>The Plant Journal</i> , 116(1): 53-62, 1998
<i>blz2</i>	endosperm	EP99106056.7
synthetic promoter	endosperm	Vicente-Carbajosa <i>et al.</i> , <i>Plant J.</i> 13: 629-640, 1998.
rice prolamin NRP33	endosperm	Wu <i>et al</i> , <i>Plant Cell Physiology</i> 39(8) 885-889, 1998
rice α -globulin Glb-1	endosperm	Wu <i>et al</i> , <i>Plant Cell Physiology</i> 39(8) 885-889, 1998
rice OSH1	embryo	Sato <i>et al</i> , <i>Proc. Natl. Acad. Sci. USA</i> , 93: 8117-8122, 1996
rice α -globulin REB/OHP-1	endosperm	Nakase <i>et al.</i> <i>Plant Mol. Biol.</i> 33: 513-522, 1997
rice ADP-glucose PP	endosperm	<i>Trans Res</i> 6:157-68, 1997
maize ESR gene family	endosperm	<i>Plant J</i> 12:235-46, 1997
sorgum γ -kafirin	endosperm	<i>PMB</i> 32:1029-35, 1996
KNOX	embryo	Postma-Haarsma <i>et al</i> , <i>Plant Mol. Biol.</i> 39:257-71, 1999
rice oleosin	embryo and aleuron	Wu <i>et at</i> , <i>J. Biochem.</i> , 123:386, 1998
sunflower oleosin	seed (embryo and dry seed)	Cummins, <i>et al.</i> , <i>Plant Mol. Biol.</i> 19: 873-876, 1992
LEAFY	shoot meristem	Weigel <i>et al.</i> , <i>Cell</i> 69:843-859, 1992.

<i>Arabidopsis thaliana</i> knat1	shoot meristem	Accession number AJ131822
<i>Malus domestica</i> kn1	shoot meristem	Accession number Z71981
<i>CLAVATA1</i>	shoot meristem	Accession number AF049870
stigma-specific genes	stigma	Nasrallah, et al., Proc. Natl. Acad. Sci. USA 85: 5551, 1988; Trick, et al., Plant Mol. Biol. 15: 203, 1990.
class I patatin gene	tuber	Liu et al., Plant Mol. Biol. 153:386-395, 1991.
PCNA rice	meristem	Kosugi et al, Nucleic Acids Research 19:1571-1576, 1991; Kosugi S. and Ohashi Y, Plant Cell 9:1607-1619, 1997.
Pea TubA1 tubulin	Dividing cells	Stotz and Long, Plant Mol.Biol. 41, 601-614. 1999
<i>Arabidopsis</i> cdc2a	cycling cells	Chung and Parish, FEBS Lett, 3;362(2):215-9, 1995
<i>Arabidopsis</i> Rop1A	Anthers; mature pollen + pollen tubes	Li et al. 1998 Plant Physiol 118, 407-417.
<i>Arabidopsis</i> AtDMC1	Meiosis-associated	Klimyuk and Jones 1997 Plant J. 11, 1-14.
Pea PS-IAA4/5 and PS-IAA6	Auxin-inducible	Wong et al. 1996 Plant J. 9, 587-599.
Pea farnesyltransferase	Meristematic tissues; phloem near growing tissues; light- and sugar-repressed	Zhou et al. 1997 Plant J. 12, 921-930
Tobacco (<i>N. sylvestris</i>) cyclin B1;1	Dividing cells / meristematic tissue	Trehin et al. 1997 Plant Mol.Biol. 35, 667-672.
<i>Catharanthus roseus</i> Mitotic cyclins CYS (A-type) and CYM (B-type)	Dividing cells / meristematic tissue	Ito et al. 1997 Plant J. 11, 983-992
<i>Arabidopsis</i> cyc1At (=cyc B1;1) and cyc3aAt (A-type)	Dividing cells / meristematic tissue	Shaul et al. 1996 Proc.Natl.Acad.Sci.U.S.A 93, 4868-4872.
<i>Arabidopsis</i> tef1 promoter box	Dividing cells / meristematic tissue	Regad et al. 1995 Mol.Gen.Genet. 248, 703-711.
<i>Catharanthus roseus</i> cyc07	Dividing cells / meristematic tissue	Ito et al. 1994 Plant Mol.Biol. 24, 863-878.

Table 4 (continued). Exemplary plant-expressible promoters for use in the performance of the present invention

II: EXEMPLARY CONSTITUTIVE PROMOTERS		
GENE SOURCE	EXPRESSION PATTERN	REFERENCE
Actin	constitutive	McElroy <i>et al</i> , Plant Cell, 2: 163-171, 1990
CAMV 35S	constitutive	Odell <i>et al</i> , Nature, 313: 810-812, 1985
CaMV 19S	constitutive	Nilsson <i>et al.</i> , Physiol. Plant. 100:456-462, 1997
GOS2	constitutive	de Pater <i>et al</i> , Plant J. 2:837-844, 1992
ubiquitin	constitutive	Christensen <i>et al</i> , Plant Mol. Biol. 18: 675-689, 1992
rice cyclophilin	constitutive	Buchholz <i>et al</i> , Plant Mol Biol. 25: 837-843, 1994
maize histone H3	constitutive	Lepetit <i>et al</i> , Mol. Gen. Genet. 231:276-285, 1992
alfalfa histone H3	constitutive	Wu <i>et al.</i> , Nucleic Acids Res. 17: 3057-3063, 1989; Wu <i>et al.</i> , Plant Mol. Biol. 11:641-649, 1988
actin 2	constitutive	An <i>et al</i> , Plant J. 10(1); 107-121, 1996

Table 4 (continued). Exemplary plant-expressible promoters for use in the performance of the present invention

III: EXEMPLARY STRESS-INDUCIBLE PROMOTERS		
NAME	STRESS	REFERENCE
P5CS (delta(1)-pyrroline-5-carboxylate syntase)	salt, water	Zhang et al. Plant Science. 129: 81-89, 1997
cor15a	cold	Hajela et al., Plant Physiol. 93: 1246-1252, 1990
cor15b	cold	Wlihelm et al., Plant Mol Biol. 23:1073-1077, 1993
cor15a (-305 to +78 nt)	cold, drought	Baker et al., Plant Mol Biol. 24: 701-713, 1994
rd29	salt, drought, cold	Kasuga et al., Nature Biotechnology 18:287-291, 1999
heat shock proteins, including artificial promoters containing the heat shock element (HSE)	heat	Barros et al., Plant Mol Biol 19: 665-75, 1992. Marrs et al., Dev Genet.14: 27-41, 1993. Schoffl et al., Mol Gen Gent, 217: 246-53, 1989.
smHSP (small heat shock proteins)	heat	Waters et al, J Experimental Botany 47:325-338, 1996
wcs120	cold	Ouellet et al., FEBS Lett. 423: 324-328, 1998
ci7	cold	Kirch et al., Plant Mol Biol 33: 897-909, 1997
Adh	cold, drought, hypoxia	Dolferus et al., Plant Physiol 105: 1075-87, 1994
pws18	water: salt and drought	Joshee et al., Plant Cell Physiol 39: 64-72, 1998
ci21A	cold	Schneider et al., Plant Physiol 113: 335-45, 1997
Trg-31	drought	Chaudhary et al., Plant Mol Biol 30: 1247-57, 1996
osmotin	osmotic	Raghothama et al., Plant Mol Biol 23: 1117-28, 1993
Rab17	osmotic, ABA	Vilardell et al., Plant Mol Biol 17: 985-93, 1991
lapA	wounding, enviromental	WO99/03977 University of California/INRA

Table 4 (continued). Exemplary plant-expressible promoters for use in the performance of the present invention

5

IV: EXEMPLARY PATHOGEN-INDUCIBLE PROMOTERS		
NAME	PATHOGEN	REFERENCE
RB7	Root-knot nematodes (Meloidogyne spp.)	US5760386 - North Carolina State University; Opperman et al (1994) <i>Science</i> 263: 221-23.
PR-1, 2, 3, 4, 5, 8, 11	fungal, viral, bacterial	Ward et al (1991) Plant Cell 3: 1085-1094; Reiss et al 1996; Lebel et al (1998), Plant J, 16(2):223-33; Melchers et al (1994), Plant J, 5(4):469-80; Lawton et al (1992), Plant Mol Biol, 19(5):735-43.
HMG2	nematodes	WO9503690 - Virginia Tech Intellectual Properties Inc .
Abi3	Cyst nematodes (Heterodera spp.)	Unpublished
ARM1	nematodes	Barthels et al., (1997) The Plant Cell 9, 2119-2134. WO 98/31822 – Plant Genetic Systems
Att0728	nematodes	Barthels et al., (1997) The Plant Cell 9, 2119-2134. <i>PCT/EP98/07761</i>
Att1712	nematodes	Barthels et al., (1997) The Plant Cell 9, 2119-2134. <i>PCT/EP98/07761</i>
Gst1	Different types of pathogens	Strittmatter et al (1996) Mol. Plant-Microbe Interact. 9, 68-73.
LEMMI	nematodes	WO 92/21757 – Plant Genetic Systems
CLE	geminivirus	<i>PCT/EP99/03445 - CINESTAV</i>
PDF1.2	Fungal including <i>Alternaria brassicicola</i> and <i>Botrytis cinerea</i>	Manners et al (1998), Plant Mol Biol, 38(6):1071-80.
Thi2.1	Fungal – <i>Fusarium oxysporum f sp. matthiolae</i>	Vignutelli et al (1998) Plant J;14(3):285-95
DB#226	nematodes	Bird and Wilson (1994) Mol. Plant-Microbe Interact., 7, 419-42 WO 95.322888

DB#280	nematodes	Bird and Wilson (1994) Mol. Plant-Microbe Interact., 7, 419-42 WO 95.322888
Cat2	nematodes	Niebel et al (1995) Mol Plant Microbe Interact 1995 May-Jun;8(3):371-8
□Tub	nematodes	Aristizabal et al (1996), 8 th International Congress on Plant-Microbe Interaction, Knoxville US B-29
SHSP	nematodes	Fenoll et al (1997) In: Cellular and molecular aspects of plant-nematode interactions. Kluwer Academic, C. Fenoll, F.M.W. Grundler and S.A. Ohi (Eds.),
Tsw12	nematodes	Fenoll et al (1997) In: Cellular and molecular aspects of plant-nematode interactions. Kluwer Academic, C. Fenoll, F.M.W. Grundler and S.A. Ohi (Eds.)
Hs1(pro1)	nematodes	WO 98/122335 - Jung
NsLTP	viral, fungal, bacterial	Molina & García-Olmedo (1993) FEBS Lett, 316(2):119-22
RIP	viral, fungal	Turner et al (1997) Proc Natl Acad Sci U S A, 94(8):3866-71

- Examples of terminators particularly suitable for use in the gene constructs of the present invention include the *Agrobacterium tumefaciens* nopaline synthase (NOS) gene terminator, the *Agrobacterium tumefaciens* octopine synthase (OCS) gene terminator sequence, the Cauliflower mosaic virus (CaMV) 35S gene terminator sequence, the *Oryza sativa* ADP-glucose pyrophosphorylase terminator sequence (t3'Bt2), the *Zea mays* zein gene terminator sequence, the *rbcS-1A* gene terminator, and the *rbcS-3A* gene terminator sequences, amongst others.
- 10 Preferred promoter sequences of the invention include root specific promoters such as but not limited to the ones listed in Table 5 and as outlined in the Examples.

Table 5. Exemplary root specific promoters for use in the performance of the present invention

NAME	ORIGIN	REFERENCE
SbPRP1	Soybean	Suzuki et al., Plant Mol Biol, 21: 109-119, 1993
636 bp fragment of TobRB7	Tobacco	Yamamoto et al., Plant Cell 3:371-382, 1991
GGPS3	Arabidopsis	Okada et al., Plant Physiol 122: 1045-1056, 2000
580 bp fragment of prxEa	Arabidopsis	Wanapu and Shinmyo, Ann N Y Acad Sci 782: 107-114, 1996
lds2 promoter	Barley	Okumura et al., Plant Mol Biol 25: 705-719, 1994
AtPRP3	Arabidopsis	Fowler et al., Plant Physiol 121: 1081-1092, 1999

5

Those skilled in the art will be aware of additional promoter sequences and terminator sequences which may be suitable for use in performing the invention. Such sequences may readily be used without any undue experimentation.

10 In the context of the current invention, "ectopic expression" or "ectopic overexpression" of a gene or a protein are conferring to expression patterns and/or expression levels of said gene or protein normally not occurring under natural conditions, more specifically is meant increased expression and/or increased expression levels. Ectopic expression can be achieved in a number of ways including operably linking of a coding sequence encoding said protein to an
15 isolated homologous or heterologous promoter in order to create a chimeric gene and/or operably linking said coding sequence to its own isolated promoter (i.e. the unisolated promoter naturally driving expression of said protein) in order to create a recombinant gene duplication or gene multiplication effect. With "ectopic co-expression" is meant the ectopic expression or ectopic overexpression of two
20 or more genes or proteins. The same or, more preferably, different promoters are used to confer ectopic expression of said genes or proteins.

Preferably, the promoter sequence used in the context of the present invention is operably linked to a coding sequence or open reading frame (ORF) encoding a cytokinin oxidase protein or a homologue, derivative or an immunologically active and/or functional fragment thereof as defined supra.

5 "Downregulation of expression" as used herein means lowering levels of gene expression and/or levels of active gene product and/or levels of gene product activity. Decreases in expression may be accomplished by e.g. the addition of coding sequences or parts thereof in a sense orientation (if resulting in co-suppression) or in an antisense orientation relative to a promoter sequence and
10 furthermore by e.g. insertion mutagenesis (e.g. T-DNA insertion or transposon insertion) or by gene silencing strategies as described by e.g. Angell and Baulcombe (1998 - WO9836083), Lowe et al. (1989 - WO9853083), Lederer et al. (1999 - WO9915682) or Wang et al. (1999 - WO9953050). Genetic constructs aimed at silencing gene expression may have the nucleotide sequence of said
15 gene (or one or more parts thereof) contained therein in a sense and/or antisense orientation relative to the promoter sequence. Another method to downregulate gene expression comprises the use of ribozymes.

Modulating, including lowering, the level of active gene products or of gene product activity can be achieved by administering or exposing cells, tissues,
20 organs or organisms to said gene product, a homologue, derivative and/or immunologically active fragment thereof. Immunomodulation is another example of a technique capable of downregulation levels of active gene product and/or of gene product activity and comprises administration of or exposing to or expressing antibodies to said gene product to or in cells, tissues, organs or
25 organisms wherein levels of said gene product and/or gene product activity are to be modulated. Such antibodies comprise "plantibodies", single chain antibodies, IgG antibodies and heavy chain camel antibodies as well as fragments thereof.

Modulating, including lowering, the level of active gene products or of gene product activity can furthermore be achieved by administering or exposing cells,
30 tissues, organs or organisms to an agonist of said gene product or the activity thereof. Such agonists include proteins (comprising e.g. kinases and

proteinases) and chemical compounds identified according to the current invention as described supra.

In the context of the current invention is envisaged the downregulation of the expression of a cytokinin oxidase gene as defined higher. Preferably said
5 cytokinin oxidase gene is a plant cytokinin oxidase gene, more specifically an *AtCKX*. The invention further comprises downregulation of levels of a cytokinin oxidase protein or of a cytokinin oxidase activity whereby said cytokinin oxidase protein has been defined supra. Preferably said cytokinin oxidase protein is a plant cytokinin oxidase, more specifically an *AtCKX*.

10 By "modifying cell fate and/or plant development and/or plant morphology and/or biochemistry and/or physiology" is meant that one or more developmental and/or morphological and/or biochemical and/or physiological characteristics of a plant is altered by the performance of one or more steps pertaining to the invention described herein.

15 "Cell fate" refers to the cell-type or cellular characteristics of a particular cell that are produced during plant development or a cellular process therefor, in particular during the cell cycle or as a consequence of a cell cycle process.

"Plant development" or the term "plant developmental characteristic" or similar term shall, when used herein, be taken to mean any cellular process of a plant
20 that is involved in determining the developmental fate of a plant cell, in particular the specific tissue or organ type into which a progenitor cell will develop. Cellular processes relevant to plant development will be known to those skilled in the art. Such processes include, for example, morphogenesis, photomorphogenesis, shoot development, root development, vegetative development, reproductive
25 development, stem elongation, flowering, and regulatory mechanisms involved in determining cell fate, in particular a process or regulatory process involving the cell cycle.

"Plant morphology" or the term "plant morphological characteristic" or similar term will, when used herein, be understood by those skilled in the art to refer to
30 the external appearance of a plant, including any one or more structural features or combination of structural features thereof. Such structural features include the

shape, size, number, position, colour, texture, arrangement, and patterning of any cell, tissue or organ or groups of cells, tissues or organs of a plant, including the root, stem, leaf, shoot, petiole, trichome, flower, petal, stigma, style, stamen, pollen, ovule, seed, embryo, endosperm, seed coat, aleurone, fibre, fruit,
5 cambium, wood, heartwood, parenchyma, aerenchyma, sieve element, phloem or vascular tissue, amongst others.

"Plant biochemistry" or the term "plant biochemical characteristic" or similar term will, when used herein, be understood by those skilled in the art to refer to the metabolic and catalytic processes of a plant, including primary and secondary
10 metabolism and the products thereof, including any small molecules, macromolecules or chemical compounds, such as but not limited to starches, sugars, proteins, peptides, enzymes, hormones, growth factors, nucleic acid molecules, celluloses, hemicelluloses, calloses, lectins, fibres, pigments such as anthocyanins, vitamins, minerals, micronutrients, or macronutrients, that are
15 produced by plants.

"Plant physiology" or the term "plant physiological characteristic" or similar term will, when used herein, be understood to refer to the functional processes of a plant, including developmental processes such as growth, expansion and differentiation, sexual development, sexual reproduction, seed set, seed
20 development, grain filling, asexual reproduction, cell division, dormancy, germination, light adaptation, photosynthesis, leaf expansion, fibre production, secondary growth or wood production, amongst others; responses of a plant to externally-applied factors such as metals, chemicals, hormones, growth factors, environment and environmental stress factors (eg. anoxia, hypoxia, high
25 temperature, low temperature, dehydration, light, daylength, flooding, salt, heavy metals, amongst others), including adaptive responses of plants to said externally-applied factors.

Means for introducing recombinant DNA into plant tissue or cells include, but are not limited to, transformation using CaCl_2 and variations thereof, in particular the
30 method described by Hanahan (1983), direct DNA uptake into protoplasts (Krens *et al*, 1982; Paszkowski *et al*, 1984), PEG-mediated uptake to protoplasts

(Armstrong *et al.*, 1990) microparticle bombardment, electroporation (Fromm *et al.*, 1985), microinjection of DNA (Crossway *et al.*, 1986), microparticle bombardment of tissue explants or cells (Christou *et al.*, 1988; Sanford, 1988), vacuum-infiltration of tissue with nucleic acid, or in the case of plants, T-DNA-mediated transfer from *Agrobacterium* to the plant tissue as described essentially
5 by An *et al.* (1985), Dodds *et al.*, (1985), Herrera-Estrella *et al.* (1983a, 1983b, 1985). Methods for transformation of monocotyledonous plants are well known in the art and include *Agrobacterium*-mediated transformation (Cheng *et al.*, 1997 - WO9748814; Hansen 1998 - WO9854961; Hiei *et al.*, 1994 - WO9400977; Hiei
10 *et al.*, 1998 - WO9817813; Rikiishi *et al.*, 1999 - WO9904618; Saito *et al.*, 1995 - WO9506722), microprojectile bombardment (Adams *et al.*, 1999 - US5969213; Bowen *et al.*, 1998 - US5736369; Chang *et al.*, 1994 - WO9413822; Lundquist *et al.*, 1999 - US5874265/US5990390; Vasil and Vasil, 1995 - US5405765. Walker *et al.*, 1999 - US5955362), DNA uptake (Eyal *et al.*, 1993 - WO9318168),
15 microinjection of *Agrobacterium* cells (von Holt, 1994 - DE4309203) and sonication (Finer *et al.*, 1997 - US5693512).

For microparticle bombardment of cells, a microparticle is propelled into a cell to produce a transformed cell. Any suitable ballistic cell transformation methodology and apparatus can be used in performing the present invention. Exemplary
20 apparatus and procedures are disclosed by Stomp *et al.* (U.S. Patent No. 5,122,466) and Sanford and Wolf (U.S. Patent No. 4,945,050). When using ballistic transformation procedures, the gene construct may incorporate a plasmid capable of replicating in the cell to be transformed.

Examples of microparticles suitable for use in such systems include 1 to 5 μm
25 gold spheres. The DNA construct may be deposited on the microparticle by any suitable technique, such as by precipitation.

A whole plant may be regenerated from the transformed or transfected cell, in accordance with procedures well known in the art. Plant tissue capable of subsequent clonal propagation, whether by organogenesis or embryogenesis,
30 may be transformed with a gene construct of the present invention and a whole plant regenerated therefrom. The particular tissue chosen will vary depending on

the clonal propagation systems available for, and best suited to, the particular species being transformed. Exemplary tissue targets include leaf disks, pollen, embryos, cotyledons, hypocotyls, megagametophytes, callus tissue, existing meristematic tissue (e.g., apical meristem, axillary buds, and root meristems),
5 and induced meristem tissue (e.g., cotyledon meristem and hypocotyl meristem).

The term "organogenesis", as used herein, means a process by which shoots and roots are developed sequentially from meristematic centres.

The term "embryogenesis", as used herein, means a process by which shoots and roots develop together in a concerted fashion (not sequentially), whether
10 from somatic cells or gametes.

Preferably, the plant is produced according to the inventive method is transfected or transformed with a genetic sequence, or amenable to the introduction of a protein, by any art-recognized means, such as microprojectile bombardment, microinjection, *Agrobacterium*-mediated transformation (including *in planta*
15 transformation), protoplast fusion, or electroporation, amongst others. Most preferably said plant is produced by *Agrobacterium*-mediated transformation.

Agrobacterium-mediated transformation or agrolistic transformation of plants, yeast, moulds or filamentous fungi is based on the transfer of part of the transformation vector sequences, called the T-DNA, to the nucleus and on
20 integration of said T-DNA in the genome of said eukaryote.

With "*Agrobacterium*" is meant a member of the *Agrobacteriaceae*, more preferably *Agrobacterium* or *Rhizobacterium* and most preferably *Agrobacterium tumefaciens*.

With "T-DNA", or transferred DNA, is meant that part of the transformation vector
25 flanked by T-DNA borders which is, after activation of the *Agrobacterium vir* genes, nicked at the T-DNA borders and is transferred as a single stranded DNA to the nucleus of an eukaryotic cell.

When used herein, with "T-DNA borders", "T-DNA border region", or "border region" are meant either right T-DNA border (RB) or left T-DNA border (LB).
30 Such a border comprises a core sequence flanked by a border inner region as part of the T-DNA flanking the border and/or a border outer region as part of the

vector backbone flanking the border. The core sequences comprise 22 bp in case of octopine-type vectors and 25 bp in case of nopaline-type vectors. The core sequences in the right border region and left border region form imperfect repeats. Border core sequences are indispensable for recognition and processing by the *Agrobacterium* nicking complex consisting of at least VirD1 and VirD2. Core sequences flanking a T-DNA are sufficient to promote transfer of said T-DNA. However, efficiency of transformation using transformation vectors carrying said T-DNA solely flanked by said core sequences is low. Border inner and outer regions are known to modulate efficiency of T-DNA transfer (Wang et al. 1987). One element enhancing T-DNA transfer has been characterized and resides in the right border outer region and is called *overdrive* (Peralta et al. 1986, van Haaren et al. 1987).

With "T-DNA transformation vector" or "T-DNA vector" is meant any vector encompassing a T-DNA sequence flanked by a right and left T-DNA border consisting of at least the right and left border core sequences, respectively, and used for transformation of any eukaryotic cell.

With "T-DNA vector backbone sequence" or "T-DNA vector backbone sequences" is meant all DNA of a T-DNA containing vector that lies outside of the T-DNA borders and, more specifically, outside the nicking sites of the border core imperfect repeats.

The current invention includes optimized T-DNA vectors such that vector backbone integration in the genome of a eukaryotic cell is minimized or absent. With "optimized T-DNA vector" is meant a T-DNA vector designed either to decrease or abolish transfer of vector backbone sequences to the genome of a eukaryotic cell. Such T-DNA vectors are known to the one familiar with the art and include those described by Hanson et al. (1999) and by Stuiver et al. (1999 - WO9901563).

The current invention clearly considers the inclusion of a DNA sequence encoding a cytokinin oxidase, homologue, derivative or immunologically active and/or functional fragment thereof as defined supra, in any T-DNA vector comprising binary transformation vectors, super-binary transformation vectors,

co-integrate transformation vectors, Ri-derived transformation vectors as well as in T-DNA carrying vectors used in agrolistic transformation. Preferably, said cytokinin oxidase is a plant cytokinin oxidase, more specifically an *Arabidopsis thaliana* (At)CKX.

5 With "binary transformation vector" is meant a T-DNA transformation vector comprising:

(a) a T-DNA region comprising at least one gene of interest and/or at least one selectable marker active in the eukaryotic cell to be transformed; and

10 (b) a vector backbone region comprising at least origins of replication active in *E. coli* and *Agrobacterium* and markers for selection in *E. coli* and *Agrobacterium*.

The T-DNA borders of a binary transformation vector can be derived from octopine-type or nopaline-type Ti plasmids or from both. The T-DNA of a binary
15 vector is only transferred to a eukaryotic cell in conjunction with a helper plasmid. With "helper plasmid" is meant a plasmid that is stably maintained in *Agrobacterium* and is at least carrying the set of *vir* genes necessary for enabling transfer of the T-DNA. Said set of *vir* genes can be derived from either octopine-type or nopaline-type Ti plasmids or from both.

20 With "super-binary transformation vector" is meant a binary transformation vector additionally carrying in the vector backbone region a *vir* region of the Ti plasmid pTiBo542 of the super-virulent *A. tumefaciens* strain A281 (EP0604662, EP0687730). Super-binary transformation vectors are used in conjunction with a helper plasmid.

25 With "co-integrate transformation vector" is meant a T-DNA vector at least comprising:

(a) a T-DNA region comprising at least one gene of interest and/or at least one selectable marker active in plants; and

30 (b) a vector backbone region comprising at least origins of replication active in *Escherichia coli* and *Agrobacterium*, and markers for selection in *E.*

coli and *Agrobacterium*, and a set of *vir* genes necessary for enabling transfer of the T-DNA.

The T-DNA borders and said set of *vir* genes of a said T-DNA vector can be derived from either octopine-type or nopaline-type Ti plasmids or from both.

5 With "Ri-derived plant transformation vector" is meant a binary transformation vector in which the T-DNA borders are derived from a Ti plasmid and said binary transformation vector being used in conjunction with a 'helper' Ri-plasmid carrying the necessary set of *vir* genes.

As used herein, the term "selectable marker gene" or "selectable marker" or
10 "marker for selection" includes any gene which confers a phenotype on a cell in which it is expressed to facilitate the identification and/or selection of cells which are transfected or transformed with a gene construct of the invention or a derivative thereof. Suitable selectable marker genes contemplated herein include the ampicillin resistance (*Amp^r*), tetracycline resistance gene (*Tc^r*), bacterial
15 kanamycin resistance gene (*Kan^r*), phosphinothricin resistance gene, neomycin phosphotransferase gene (*nptII*), hygromycin resistance gene, β -glucuronidase (GUS) gene, chloramphenicol acetyltransferase (CAT) gene, green fluorescent protein (*gfp*) gene (Haseloff *et al*, 1997), and luciferase gene, amongst others.

With "agrolistics", "agrolistic transformation" or "agrolistic transfer" is meant here
20 a transformation method combining features of *Agrobacterium*-mediated transformation and of biolistic DNA delivery. As such, a T-DNA containing target plasmid is co-delivered with DNA/RNA enabling in planta production of VirD1 and VirD2 with or without VirE2 (Hansen and Chilton 1996; Hansen *et al*. 1997; Hansen and Chilton 1997 - WO9712046).

25 With "foreign DNA" is meant any DNA sequence that is introduced in the host's genome by recombinant techniques. Said foreign DNA includes e.g. a T-DNA sequence or a part thereof such as the T-DNA sequence comprising the selectable marker in an expressible format. Foreign DNA furthermore include intervening DNA sequences as defined supra.

30 With "recombination event" is meant either a site-specific recombination event or a recombination event effected by transposon 'jumping'.

With "recombinase" is meant either a site-specific recombinase or a transposase. With "recombination site" is meant either site-specific recombination sites or transposon border sequences.

With "site specific recombination event" is meant an event catalyzed by a system
5 generally consisting of three elements: a pair of DNA sequences (the site-specific recombination sequences or sites) and a specific enzyme (the site-specific recombinase). The site-specific recombinase catalyzes a recombination reaction only between two site-specific recombination sequences depending on the orientation of the site-specific recombination sequences. Sequences intervening
10 between two site-specific recombination sites will be inverted in the presence of the site-specific recombinase when the site-specific recombination sequences are oriented in opposite directions relative to one another (i.e. inverted repeats). If the site-specific recombination sequences are oriented in the same direction relative to one another (i.e. direct repeats), then any intervening sequences will
15 be deleted upon interaction with the site-specific recombinase. Thus, if the site-specific recombination sequences are present as direct repeats at both ends of a foreign DNA sequence integrated into a eukaryotic genome, such integration of said sequences can subsequently be reversed by interaction of the site-specific recombination sequences with the corresponding site specific recombinase.

20 A number of different site specific recombinase systems can be used including but not limited to the Cre/lox system of bacteriophage P1, the FLP/FRT system of yeast, the Gin recombinase of phage Mu, the Pin recombinase of *E. coli*, the PinB, PinD and PinF from *Shigella*, and the R/RS system of the pSR1 plasmid. Recombinases generally are integrases, resolvases or flippases. Also dual-
25 specific recombinases can be used in conjunction with direct or indirect repeats of two different site-specific recombination sites corresponding to the dual-specific recombinase (WO99/25840). The two preferred site-specific recombinase systems are the bacteriophage P1 Cre/lox and the yeast FLP/FRT systems. In these systems a recombinase (Cre or FLP) interact specifically with
30 its respective site-specific recombination sequence (lox or FRT respectively) to invert or excise the intervening sequences. The site-specific recombination

sequences for each of these two systems are relatively short (34 bp for lox and 47 bp for FRT). Some of these systems have already been used with high efficiency in plants such as tobacco (Dale et al. 1990) and *Arabidopsis* (Osborne et al. 1995). Site-specific recombination systems have many applications in plant molecular biology including methods for control of homologous recombination (e.g. US5527695), for targeted insertion, gene stacking, etc. (WO99/25821) and for resolution of complex T-DNA integration patterns or for excision of a selectable marker (WO99/23202).

Although the site-specific recombination sequences must be linked to the ends of the DNA to be excised or to be inverted, the gene encoding the site specific recombinase may be located elsewhere. For example, the recombinase gene could already be present in the eukaryote's DNA or could be supplied by a later introduced DNA fragment either introduced directly into cells, through crossing or through cross-pollination. Alternatively, a substantially purified recombinase protein could be introduced directly into the eukaryotic cell, e.g. by micro-injection or particle bombardment. Typically, the site-specific recombinase coding region will be operably linked to regulatory sequences enabling expression of the site-specific recombinase in the eukaryotic cell.

With "recombination event effected by transposon jumping" or "transposase-mediated recombination" is meant a recombination event catalyzed by a system consisting of three elements: a pair of DNA sequences (the transposon border sequences) and a specific enzyme (the transposase). The transposase catalyzes a recombination reaction only between two transposon border sequences which are arranged as inverted repeats.

A number of different transposon/transposase systems can be used including but not limited to the Ds/Ac system, the Spm system and the Mu system. These systems originate from corn but it has been shown that at least the Ds/Ac and the Spm system also function in other plants (Fedoroff et al. 1993, Schlappi et al. 1993, Van Sluys et al. 1987). Preferred are the Ds- and the Spm-type transposons which are delineated by 11 bp- and 13 bp- border sequences, respectively.

Although the transposon border sequences must be linked to the ends of the DNA to be excised, the gene encoding the transposase may be located elsewhere. For example, the recombinase gene could already be present in the eukaryote's DNA or could be supplied by a later introduced DNA fragment either
5 introduced directly into cells, through crossing or through cross-pollination. Alternatively, a substantially purified transposase protein could be introduced directly into cells, e.g. by microinjection or by particle bombardment.

As part of the current invention, transposon border sequences are included in a foreign DNA sequence such that they lie outside said DNA sequence and
10 transform said DNA into a transposon-like entity that can move by the action of a transposase.

As transposons often reintegrate at another locus of the host's genome, segregation of the progeny of the hosts in which the transposase was allowed to act might be necessary to separate transformed hosts containing e.g. only the
15 transposon footprint and transformed hosts still containing the foreign DNA.

In performing the present invention, the genetic element is preferably induced to mobilize, such as, for example, by the expression of a recombinase protein in the cell which contacts the integration site of the genetic element and facilitates a recombination event therein, excising the genetic element completely, or
20 alternatively, leaving a "footprint", generally of about 20 nucleotides in length or greater, at the original integration site. Those hosts and host parts that have been produced according to the inventive method can be identified by standard nucleic acid hybridization and/or amplification techniques to detect the presence of the mobilizable genetic element or a gene construct comprising the same.
25 Alternatively, in the case of transformed host cells, tissues, and hosts wherein the mobilizable genetic element has been excised, it is possible to detect a footprint in the genome of the host which has been left following the excision event, using such techniques. As used herein, the term "footprint" shall be taken to refer to any derivative of a mobilizable genetic element or gene construct
30 comprising the same as described herein which is produced by excision, deletion or other removal of the mobilizable genetic element from the genome of a cell

transformed previously with said gene construct. A footprint generally comprises at least a single copy of the recombination loci or transposon used to promote excision. However, a footprint may comprise additional sequences derived from the gene construct, for example nucleotide sequences derived from the left border sequence, right border sequence, origin of replication, recombinase-
5 encoding or transposase-encoding sequence if used, or other vector-derived nucleotide sequences. Accordingly, a footprint is identifiable according to the nucleotide sequence of the recombination locus or transposon of the gene construct used, such as, for example, a sequence of nucleotides corresponding
10 or complementary to a *lox* site or *frt* site.

The term "cell cycle" means the cyclic biochemical and structural events associated with growth and with division of cells, and in particular with the regulation of the replication of DNA and mitosis. Cell cycle includes phases called: G0, Gap1 (G1), DNA synthesis (S), Gap2 (G2), and mitosis (M). Normally
15 these four phases occur sequentially, however, the cell cycle also includes modified cycles wherein one or more phases are absent resulting in modified cell cycle such as endomitosis, acytokinesis, polyploidy, polyteny, and endoreduplication.

The term "cell cycle progression" refers to the process of passing through the different cell cycle phases. The term "cell cycle progression rate" accordingly
20 refers to the speed at which said cell cycle phases are run through or the time spans required to complete said cell cycle phases.

With "two-hybrid assay" is meant an assay that is based on the observation that many eukaryotic transcription factors comprise two domains, a DNA-binding
25 domain (DB) and an activation domain (AD) which, when physically separated (i.e. disruption of the covalent linkage) do not effectuate target gene expression. Two proteins able to interact physically with one of said proteins fused to DB and the other of said proteins fused to AD will re-unite the DB and AD domains of the transcription factor resulting in target gene expression. The target gene in the
30 yeast two-hybrid assay is usually a reporter gene such as the β -galactosidase gene. Interaction between protein partners in the yeast two-hybrid assay can

thus be quantified by measuring the activity of the reporter gene product (Bartel and Fields 1997). Alternatively, a mammalian two-hybrid system can be used which includes e.g. a chimeric green fluorescent protein encoding reporter gene (Shioda *et al.*, 2000).

5 Furthermore, folding simulations and computer redesign of structural motifs of the protein of the invention can be performed using appropriate computer programs (Olszewski, *Proteins* 25 (1996), 286-299; Hoffman, *Comput. Appl. Biosci.* 1 (1995), 675-679). Computer modeling of protein folding can be used for the conformational and energetic analysis of detailed peptide and protein models
10 (Monge, *J. Mol. Biol.* 247 (1995), 995-1012; Renouf, *Adv. Exp. Med. Biol.* 376 (1995), 37-45). In particular, the appropriate programs can be used for the identification of interactive sites of the cytokinin oxidases, its ligands or other interacting proteins by computer assistant searches for complementary peptide sequences (Fassina, *Immunomethods* 5 (1994), 114-120). Further appropriate
15 computer systems for the design of protein and peptides are described in the prior art, for example in Berry, *Biochem. Soc. Trans.* 22 (1994), 1033-1036; Wodak, *Ann. N. Y. Acad. Sci.* 501 (1987), 1-13; Pabo, *Biochemistry* 25 (1986), 5987-5991. The results obtained from the above-described computer analysis can be used for, e.g. the preparation of peptidomimetics of the protein of the
20 invention or fragments thereof. Such pseudopeptide analogues of the natural amino acid sequence of the protein may very efficiently mimic the parent protein (Benkirane, *J. Biol. Chem.* 271 (1996), 33218-33224). For example, incorporation of easily available achiral Ω -amino acid residues into a protein of the invention or a fragment thereof results in the substitution of amino bonds by polymethylene
25 units of an aliphatic chain, thereby providing a convenient strategy for constructing a peptidomimetic (Banerjee, *Biopolymers* 39 (1996), 769-777). Superactive peptidomimetic analogues of small peptide hormones in other systems are described in the prior art (Zhang, *Biochem. Biophys. Res. Commun.* 224 (1996), 327-331). Appropriate peptidomimetics of the protein of the present
30 invention can also be identified by the synthesis of peptidomimetic combinatorial libraries through successive amine alkylation and testing the resulting

compounds, e.g., for their binding, kinase inhibitory and/or immunological properties. Methods for the generation and use of peptidomimetic combinatorial libraries are described in the prior art, for example in Ostresh, Methods in Enzymology 267 (1996), 220-234 and Dorner, Bioorg. Med. Chem. 4 (1996),
5 709-715.

Furthermore, a three-dimensional and/or crystallographic structure of the protein of the invention can be used for the design of peptidomimetic inhibitors of the biological activity of the protein of the invention (Rose, Biochemistry 35 (1996), 12933-12944; Ruterber, Bioorg. Med. Chem. 4 (1996), 1545-1558).

10 The compounds to be obtained or identified in the methods of the invention can be compounds that are able to bind to any of the nucleic acids, peptides or proteins of the invention. Other interesting compounds to be identified are compounds that modulate the expression of the genes or the proteins of the invention in such a way that either the expression of said gene or protein is
15 enhanced or decreased by the action of said compound. Alternatively the compound can exert his action by enhancing or decreasing the activity of any of the proteins of the invention. Herein, preferred proteins are novel cytokinin oxidases.

Said compound or plurality of compounds may be comprised in, for example,
20 samples, e.g., cell extracts from, e.g., plants, animals or microorganisms. Furthermore, said compound(s) may be known in the art but hitherto not known to be capable of suppressing or activating cell cycle interacting proteins. The reaction mixture may be a cell free extract of may comprise a cell or tissue culture. Suitable set ups for the method of the invention are known to the person
25 skilled in the art and are, for example, generally described in Alberts et al., Molecular Biology of the Cell, third edition (1994), in particular Chapter 17. The plurality of compounds may be, e.g., added to the reaction mixture, culture medium or injected into the cell.

If a sample containing a compound or a plurality of compounds is identified in the
30 method of the invention, then it is either possible to isolate the compound form the original sample identified as containing the compound capable of acting as

an agonist, or one can further subdivide the original sample, for example, if it consists of a plurality of different compounds, so as to reduce the number of different substances per sample and repeat the method with the subdivisions of the original sample. Depending on the complexity of the samples, the steps
5 described above can be performed several times, preferably until the sample identified according to the method of the invention only comprises a limited number of or only one substance(s). Preferably said sample comprises substances or similar chemical and/or physical properties, and most preferably said substances are identical. Preferably, the compound identified according to
10 the above described method or its derivative is further formulated in a form suitable for the application in plant breeding or plant cell and tissue culture.

The term "early vigor" refers to the ability of a plant to grow rapidly during early development, and relates to the successful establishment, after germination, of a well-developed root system and a well-developed photosynthetic apparatus.

15 The term "lodging" refers to the ability of a plant to fix itself to the soil, and relates to the size, depth and morphology of the root system.

The following examples and figures are given by means of illustration of the present invention and are in no way limiting. The contents of all references included in this application are incorporated by reference.

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1. Schematic representation of plant cytokinin oxidase genes.

Shown are the structures of different cytokinin oxidase genes isolated from maize
5 (ZmCKX1, accession number AF044603, Biochem. Biophys. Res. Com. 255:328-
333, 1999) and Arabidopsis (AtCKX1 to AtCKX4). Exons are denominated with
'E' and represented by shaded boxes. Introns are represented by white boxes.
Further indicated are the gene sizes (in kb, on top of each structure), the gene
accession numbers (under the names) and a size bar representing 0.5 kb.

10

Figure 2. Alignment of plant cytokinin oxidase amino acid sequences.

The amino acid sequences from cytokinin oxidases from maize (ZmCKX1) and
Arabidopsis (AtCKX1 to AtCKX4) are aligned. Identical amino acid residues
identical are marked by a black box, similar amino acid residues are in a grey
15 box. Amino acid similarity groups: (M,I,L,V), (F,W,Y), (G,A), (S,T), (R,K,H), (E,D),
(N,Q),

Figure 3. Northern blot analysis of *AtCKX1*-expressing tobacco and *Arabidopsis* plants.

20 **(A)** Northern blot analysis of constitutively expressing tobacco plants (lanes 1-8)
compared to wild type SNN tobacco (lane 9)

(B) Comparison of tetracycline-induced gene expression in leaves after 12h of
induction with a constitutively-expressing clone. Lanes 2-9, leaves of four
different *CKX1*-W38TetR clones (+, -, with or without tetracycline treatment), lane
25 1, constitutively expressing 35S::*CKX1* clone.

(C) Northern blot analysis of *Arabidopsis* plants constitutively expressing *CKX1*
gene. Lanes 2-4, three different constitutively expressing 35S::*CKX1* clones
compared to wild type *Arabidopsis* plant (lane 1).

30

Figure 4: Growth characteristics of 35S::CKX1 transgenic *Arabidopsis* plants.

- (A) Two wild type seedlings (left) compared to two 35S::CKX1 expressing seedlings (right). Note the increase formation of adventitious roots and increased root branching in the transgenic seedlings. Pictures were taken 14 days after germination. Plants were grown *in vitro* on MS medium in petri dishes in a vertical position.
- (B) Like A, but roots stained with toluidine blue.
- (C) Top view of a petri dish with 35S::CKX1 transgenic seedlings three weeks after germination.
- (D) A 35S::CKX1 transgenic plants grown in liquid culture. Roots of wild type seedlings grow poorly under these conditions (not shown).
- (E) Transformants (T0) that express the 35S::CKX1 gene (three plants on the right), a wild type plant is shown on the left.
- (F) Phenotype of T1 plants grown in soil. Wild type plant (left) compared to two 35S::CKX1 transgenic plants.

Figure 5: 35S::CKX1 expression in tobacco plants.

- (A) 35S::CKX1 expressing plant (left) and wild type plant (tobacco SNN).
- (B) Close-up of the 35S::CKX1 transgenic plants. Note the growth of axillary shoot meristems.
- (C) Close-up of a strong expresser of CKX1.
- (D) Root system of 35S::CKX1 transgenic plant (left) and wild type (tobacco Wisconsin 38).

Figure 6. Northern blot analysis of AtCKX2-expressing tobacco and *Arabidopsis* plants.

- (A) Northern blot analysis of constitutively expressing tobacco plants (lanes 1-7) compared to wild type SNN tobacco (lane 8)

(B) Northern blot analysis of *Arabidopsis* plants constitutively expressing *CKX2* gene. Lanes 2-8, seven different constitutively expressing *35S::CKX1* clones compared to wild-type *Arabidopsis* plant (lane 1).

5 **Figure 7: Phenotype of *CKX2* overexpressing plants.**

T1 generation of *35S::CKX2* expressing *Arabidopsis* plants (two plants on the right) compared to wild type (plant on the left).

Figure 8: Phenotype of *CKX2* overexpressing plants.

- 10 *35S::CKX2* expressing tobacco plants (two plants on the right) compared to wild type (tobacco SNN plant on the left).

EXAMPLES

Example 1. Identification of candidate cytokinin oxidase encoding genes from *Arabidopsis thaliana*

5 Six different genes are identified from *Arabidopsis thaliana* that bear sequence similarity to a cytokinin oxidase gene from maize (Morris *et al.*, Biochem Biophys Res Comm 255:328-333, 1999; Houda-Herin *et al.* Plant J 17:615-626; WO 99/06571). These genes are found by screening 6-frame translations of nucleotide sequences from public genomic databases with the maize protein
10 sequence, employing tblastn program. These sequences are designated as *Arabidopsis thaliana* cytokinin oxidase-like genes or *AtCKX*. They are arbitrarily numbered as *AtCKX1* to *AtCKX6*. The below list summarizes the information on these genes. The predicted ORF borders and protein sequences are indicative, in order to illustrate by approximation the protein sequence divergence between
15 the *Arabidopsis* and maize cytokinin oxidases, as well as amongst the different *Arabidopsis* cytokinin oxidases. The ORF borders and protein sequences shown should not be taken as conclusive evidence for the mode of action of these *AtCKX* genes. The % homology/similarity values that are indicated amongst the *Arabidopsis* genes represent respectively the lowest and highest values found
20 with all combinations.

A. Gene name: *AtCKX1* (*Arabidopsis thaliana* cytokinin oxidase-like protein 1, SEQ ID NO1)

25 **Location in database** (accession number, location on bac): AC002510, *Arabidopsis thaliana* chromosome II section 225 of 255 of the complete sequence. Sequence from clones T32G6.

ORF predicted in the database:

30 15517..16183, 16415..16542, 16631..16891, 16995..17257, 17344..17752

Predicted protein sequence: SEQ ID NO 2

Homologies

% homology with *Z. mays* gene:

31,5% (Dnastar/MegAlign - Clustal method)

5

% similarity with *Z. mays* protein:

32,2% (Dnastar/MegAlign - Clustal method)

% homologies with other *Arabidopsis* genes (range):

10 38,2% (*AtCKX2*) - 54,1% (*AtCKX6*) (Dnastar/MegAlign - Clustal method)

% similarity with other *Arabidopsis* proteins (range):

37,1% (*AtCKX2*) - 58,1% (*AtCKX6*) (Dnastar/MegAlign - Clustal method)

15 **B. Gene name:** *AtCKX2* (*Arabidopsis thaliana* cytokinin oxidase-like protein 2, SEQ ID NO3)

Location in database (accession number, location on bac): AC005917, *Arabidopsis thaliana* chromosome II section 113 of 255 of the complete
20 sequence. Sequence from clones F27F23, F3P11.

ORF predicted in the database:

complement, 40721..41012, 41054..41364, 41513..41770, 42535..42662,
43153..43711

25

Please note: The cDNA sequence is different than annotated in the database. Based on the cDNA sequence and using the gene prediction program (NetPlantGene, <http://www.cbs.dtu.dk/services/NetGene2/>) the ORF predicted in the database is revised:

30 complement, 40721..41012, 41095..41364, 41513..41770, 42535..42662, 43153..43711

Predicted protein sequence, based on corrected ORF: SEQ ID NO 4

Homologies

% homology with *Z. mays* gene:

38,4% (Dnastar/MegAlign - Clustal method)

5 % similarity with *Z. mays* protein:

37,5% (Dnastar/MegAlign - Clustal method)

% homologies with other *Arabidopsis* genes (range):

34,9% (*AtCKX6*) – 64,5% (*AtCKX4*) (Dnastar/MegAlign - Clustal method)

10

% similarity with other *Arabidopsis* proteins (range):

36,5% (*AtCKX6*) – 66,1% (*AtCKX4*) (Dnastar/MegAlign - Clustal method)

15 **C. Gene name: *AtCKX3*** (*Arabidopsis thaliana* cytokinin oxidase-like protein 3, SEQ ID NO 5)

Location in database (accession number, location on bac): AB024035, *Arabidopsis thaliana* genomic DNA, chromosome 5, P1 clone: MHM17, complete sequence.

20

No prediction of the ORF in the database.

The gene is identified by several gene prediction programs including GRAIL (<ftp://arthur.epm.ornl.gov/pub/xgrail>), Genscan (<http://CCR-081.mit.edu/GENSCAN.html>) and (NetPlantGene, <http://www.cbs.dtu.dk/services/NetGene2/>):

25

complement, 29415..29718, 29813..30081, 30183..30443, 30529..30656, 32107..32716

Predicted protein sequence, based on own ORF prediction: SEQ ID NO 6

30

Homologies

% homology with *Z. mays* gene:

38,7% (Dnastar/MegAlign - Clustal method)

% similarity with *Z. mays* protein:

39,2% (Dnastar/MegAlign - Clustal method)

% homologies with other *Arabidopsis* genes (range):

5 38,8% (*AtCKX6*) – 51,0% (*AtCKX2*) (Dnastar/MegAlign - Clustal method)

% similarity with other *Arabidopsis* proteins (range):

39,9% (*AtCKX6*) – 46,7% (*AtCKX2*) (Dnastar/MegAlign - Clustal method)

10 **D. Gene name:** *AtCKX4* (*Arabidopsis thaliana* cytokinin oxidase-like protein 4,
SEQ ID NO 7)

Location in database (accession number, location on bac):

1) AL079344, *Arabidopsis thaliana* DNA chromosome 4, BAC clone T16L4
15 (ESSA project)

2) AL161575, *Arabidopsis thaliana* DNA chromosome 4, contig fragment No. 71.

ORF predicted in the database:

1) 76187..76814, 77189..77316, 77823..78080, 78318..78586, 78677..78968
20 2) 101002..101629, 102004..102131, 102638..102895, 103133..103401,
103492..103783

Predicted protein sequence: SEQ ID NO 8

25 **Homologies**

% homology with *Z. mays* gene:

41,0% (Dnastar/MegAlign - Clustal method)

% similarity with *Z. mays* protein:

30 41,0% (Dnastar/MegAlign - Clustal method)

% homologies with other *Arabidopsis* genes (range):

35,2% (*AtCKX6*) – 64,5% (*AtCKX2*) (Dnastar/MegAlign - Clustal method)

% similarity with other *Arabidopsis* proteins (range):

35,1% (*AtCKX6*) – 66,1% (*AtCKX2*) (Dnastar/MegAlign - Clustal method)

5 **E. Gene name:** *AtCKX5* (*Arabidopsis thaliana* cytokinin oxidase-like protein 5,
SEQ ID NO 9)

Location in database (accession number, location on bac): AC023754, F1B16,
complete sequence, chromosome 1

10 **No prediction of the ORF in the database.**

The gene can be identified by several gene prediction programs including GRAIL .
(<ftp://arthur.epm.ornl.gov/pub/xgrail>), Genscan (<http://CCR-081.mit.edu/GENSCAN.html>) and (NetPlantGene, <http://www.cbs.dtu.dk/services/NetGene2/>)

15 43756..44347, 44435..44562, 44700..44966, 45493..45755, 46200..46560

Predicted protein sequence: SEQ ID NO 10

Homologies

20 % homology with *Z. mays* gene:

39,1% (Dnastar/MegAlign - Clustal method)

% similarity with *Z. mays* protein:

36,6% (Dnastar/MegAlign - Clustal method)

25 % homologies with other *Arabidopsis* genes (range):

40,1% (*AtCKX2*) – 44,0% (*AtCKX3*) (Dnastar/MegAlign - Clustal method)

% similarity with other *Arabidopsis* proteins (range):

30 41,6% (*AtCKX4*) – 46,4% (*AtCKX6*) (Dnastar/MegAlign - Clustal method)

F. Gene name: *AtCKX6* (*Arabidopsis thaliana* cytokinin oxidase-like protein 6, SEQ ID NO 11)

Location in database (accession number, location on bac): AL163818,
5 *Arabidopsis thaliana* DNA chromosome 3, P1 clone MAA21 (ESSA project).

ORF predicted in the database:

46630..47215, 47343..47470, 47591..47806, 47899..48161, 48244..48565

10 **Predicted protein sequence:** SEQ ID NO 12

Homologies

% homology with *Z. mays* gene:

37,3% (Dnastar/MegAlign - Clustal method)

15

% similarity with *Z. mays* protein:

36,1% (Dnastar/MegAlign - Clustal method)

% homologies with other *Arabidopsis* genes (range):

20

34,9% (*AtCKX2*) – 54,1% (*AtCKX1*) (Dnastar/MegAlign - Clustal method)

% similarity with other *Arabidopsis* proteins (range):

35,1% (*AtCKX4*) – 58,1% (*AtCKX1*) (Dnastar/MegAlign - Clustal method)

25 Genes *AtCKX3* and *AtCKX5* are not annotated as putative cytokinin oxidases in the database and ORFs for these genes are not given. Furthermore, the ORF (and consequently the protein structures) predicted for *AtCKX2* is different from our own prediction.

A comparison of the gene structure of the *Arabidopsis AtCKX* genes 1 to 4 and
30 the maize *CKX* gene is shown in Fig 1.

The predicted proteins encoded by the *Arabidopsis AtCKX* genes show between 32% and 41% sequence similarity with the maize protein, while they show between 35% and 66% sequence similarity to each other. Because of this

reduced sequence conservation, it is not clear *a priori* whether the *Arabidopsis AtCKX* genes encode proteins with cytokinin oxidase activity. An alignment of the *Arabidopsis AtCKX* predicted proteins 1 to 4 and the maize *CKX* gene is shown in Fig 2 .

5

Example 2. Transgenic plants overexpressing AtCKX1 show increased cytokinin oxidase activity and altered plant morphology

1. Description of the cloning process

10 The following primers are used to PCR amplify the AtCKX1 gene from *Arabidopsis thaliana*, accession Columbia:

Sequence of 5' primer: cggtcgacATGGGATTGACCTCATCCTTACG (SEQ ID NO:13)

15 Sequence of 3' primer: gcgtcgacTTATACAGTTCTAGGTTTCGGCAGTAT (SEQ ID NO: 14)

A 2235-bp PCR fragment, amplified by these primers, is inserted in the Sal I site of pUC19. The insert is sequenced to confirm that the PCR amplification has not introduced mutations. The Sall/Sall fragment of this vector is subcloned in the
20 Sall site downstream of a modified CaMV 35S promoter (carrying three tetracycline operator sequences) in the binary vector pBinHyg-Tx (Gatz *et al.*, 1992).

The resulting construct is introduced into tobacco and *Arabidopsis thaliana* through *Agrobacterium*-mediated transformation, using standard transformation
25 protocols.

2. Molecular analysis of the transgenic lines

Several transgenic lines are identified that synthesize the *AtCKX1* transcript at high levels (Fig 3). Transgenic lines expressing *AtCKX1* transcript also show
30 increased cytokinin oxidase activity. This is exemplified for 2 tobacco and 2

Arabidopsis lines in Table 6. This result proves that the *AtCKX1* gene encodes a protein with cytokinin oxidase activity.

Table 6. Cytokinin oxidase activity in *AtCKX1* transgenic plant tissues

5

Leaf sample		
Plant species	Plant line	Cytokinin oxidase activity (nmol Ade/mg protein.h)
<i>Arabidopsis</i>	Col-0 wild-type	0.009
	CKX1-11	0.024
	CKX1-22	0.026
	CKX1-22	0.027
Tobacco	SNN wild-type	0.004
	CKX1-SNN-8	0.016
	CKX1-SNN-28	0.021

3. Phenotypic description of the transgenic lines

In *Arabidopsis* (see also Fig 4):

- 10
- onset of germination same as for wt
 - the number of adventitious roots early after germination is enhanced; the growth of aerial organs (hypocotyl, cotyledons and leaves) is retarded, dwarfed phenotype
 - the number of leaves in rosette is increased
- 15
- the life cycle is longer compared to wt

In tobacco (see also Fig 5):

The plants have a dwarfed phenotype, with reduced apical dominance and increased root production.

20

Five categories of phenotype:

- 1) strong - 2 clones
- 2) intermediate - 3 clones

- 3) weak - 4 clones
- 4) tall plants (as WT) with large inflorescence - 5 clones_g
- 5) similar to wt, 9 clones

5 Height

- wt: 90-100 cm
- weak: app. 75 cm
- intermediate: app. 40-45 cm (main stem app. 25 cm but overgrown by side branches.
- 10 - strong: app. 10 cm

Roots

Plants highly expressing the gene are easily distinguishable in vitro by their ability to form more roots which are thicker (stronger) as well as by forming aerial roots along the stem.

15

Roots in soil (4-4,5 month after transfer to the green house):

- strong: good root system (as WT) despite the strong reduction of aerial part
- intermediate: large increase in root system (biomass)
- 20 - weak: also increase in root biomass compared to WT

Internodes distance

- wt: 5th internode below inflorescence app. 5 cm, 9th internode about 2 cm long
- 25 - intermediate class: 5th about 2.5 cm, 9th internode app. 0,5 cm long

Seed capsules

Expressing plants have in general less capsules, which are of similar size as wt (even in strong expressors)

30

- strong: 2-4 capsules
- intermediate: about 25

- wt: more than 60

Flower

Size is not reduced. The first flowers are aborted in the transgenic expressers..

5

Leaves

The shape of leaves of transgenic plants (expressors) is lanceolate (longer and narrow). Basal leaves of strongly expressing plants are darker green.

The texture of older leaves is more more rigid.

10 They become yellow in the intercostal regions but stay dark green along the veins.

Stem

More side branches are formed (reduction of apical dominance). The side

15 branches overgrow the main stem (it gives the height of 40-45 cm for intermediate plants). Even secondary branches appear.

Example 3. Transgenic plants overexpressing AtCKX2 show increased cytokinin oxidase activity and altered plant morphology

20

1. Description of the cloning process

The following primers are used to PCR amplify the AtCKX2 gene from *Arabidopsis thaliana*, accession Columbia:

25 Sequence of 5' primer: gcggtaccAGAGAGAGAAACATAAACAAATGGC (SEQ ID NO:15)

Sequence of 3' primer: gcggtaccCAATTTTACTTCCACCAAATGC (SEQ ID NO:16)

30 A 3104-bp PCR fragment, amplified by these primers, is inserted in the KpnI site of pUC19. The insert is sequenced to check that no differences to the published sequence are introduced by the PCR procedure. The KpnI/KpnI fragment of this vector is subcloned in the KpnI site downstream of a modified CaMV 35S

promoter (carrying three tetracycline operator sequences) in the binary vector pBinHyg-Tx (Gatz *et al.*, 1992).

The resulting construct is introduced into tobacco and *Arabidopsis thaliana* through *Agrobacterium*-mediated transformation, using standard transformation protocols.

2. Molecular analysis of the transgenic lines

Several transgenic lines are identified that synthesize the *AtCKX2* transcript at high levels (Fig 6). Transgenic lines expressing *AtCKX2* transcript also show increased cytokinin oxidase activity. This is exemplified for 2 tobacco and 3 *Arabidopsis* lines in Table 7. This result proves that the *AtCKX2* gene encodes a protein with cytokinin oxidase activity.

Table 7. Cytokinin oxidase activity in *AtCKX2* transgenic plant tissues

Sample		
Plant species and tissue	Plant line	Cytokinin oxidase activity (nmol Ade/mg protein.h)
<i>Arabidopsis</i> callus	Col-0 wild-type	0.037
	CKX2-15	0.351
	CKX2-17	0.380
	CKX2-55	0.265
Tobacco leaves	SNN wild-type	0.009
	CKX2-SNN-18	0.091
	CKX2-SNN-19	0.091

3. Phenotypic description of the transgenic lines

In *Arabidopsis* (see Fig 7):

Basically similar to *CKX1* (no quantitative comparisons with *CKX1* available)

In tobacco (see Fig 8):

Three categories of phenotype:

- 1) strong - 15 clones (similar to intermediate phenotype of *CKX1*)
- 5 2) weak - 6 clones
- 3) others - similar to wt plants, 7 clones

Roots

Plants highly expressing the gene are easily distinguishable in vitro by their
10 ability to form more roots which are thicker (stronger) as well as by forming aerial
roots along the stem.

Roots in soil (4-4,5 month after transfer to the green house):

- strong: very significant increase in root biomass for strongest
expressing clones
- 15 - weak: also increase in root biomass compared to wt

The observations concerning plant height, internode distance, branching, seed
set, leaf form and yellowing are similar as for *AtCKX1* transgenics with some
generally minor quantitative differences.

20 Interestingly, the increase in cytokinin oxidase activity and in root production is
more pronounced in *AtCKX2* than in *AtCKX1* expressing tobacco lines, indicating
that *AtCKX2* may be a preferential tool for the embodiments of this patent
application as compared to *AtCKX1*.

25 **Example 4. Transgenic plants overexpressing AtCKX3 show altered plant
morphology**

1. Description of the cloning process

The following primers are used to PCR amplify the *AtCKX3* gene from

30 *Arabidopsis thaliana*, accession Columbia:

Sequence of 5' primer: gcggtaccTTCATTGATAAGAATCAAGCTATTCA (SEQ ID
NO:17)

Sequence of 3' primer: gcggtaccCAAAGTGGTGAGAACGACTAACA (SEQ ID NO:18)

5 A 3397-bp PCR fragment, produced by this PCR amplification, is inserted in the KpnI site of pBluescript. The insert is sequenced to confirm that the PCR product has no sequence changes as compared to the gene. The KpnI/KpnI fragment of this vector is subcloned in the KpnI site downstream of a modified CaMV 35S promoter (carrying three tetracycline operator sequences) in the binary vector pBinHyg-Tx (Gatz *et al.*, 1992).

10

2. Plant phenotypic analysis

The phenotypes generated by overexpression of the *AtCKX3* gene in tobacco and *Arabidopsis* are basically similar as those of *AtCKX1* and *AtCKX2* expressing plants, i.e. enhanced rooting and dwarfing.

15

Example 5. Transgenic plants overexpressing AtCKX4 show altered plant morphology

1. Description of the cloning process

20 The following primers are used to PCR amplify the *AtCKX4* gene from *Arabidopsis thaliana*, accession Columbia:

Sequence of 5' primer: gcggtaccCCCATTAACCTACCCGTTTG (SEQ ID NO:19)

Sequence of 3' primer: gcggtaccAGACGATGAACGTACTTGTCTGTA (SEQ ID NO:20)

25

A 2890-bp PCR fragment, produced by this PCR amplification, is inserted in the KpnI site of pBluescript. The insert is sequenced to confirm that the PCR product has no sequence changes as compared to the gene. The KpnI/KpnI fragment of this vector is subcloned in the KpnI site downstream of a modified CaMV 35S promoter (carrying three tetracycline operator sequences) in the binary vector pBinHyg-Tx (Gatz *et al.*, 1992).

30

2. Molecular analysis of the transgenic lines

Several transgenic lines are identified that show increased cytokinin oxidase activity. This is exemplified for 3 *Arabidopsis* lines in Table 8. This result proves
5 that the *AtCKX4* gene encodes a protein with cytokinin oxidase activity.

Table 8. Cytokinin oxidase activity in *AtCKX4* transgenic plant tissues

Sample		
Plant species and tissue	Plant line	Cytokinin oxidase activity (nmol Ade/mg protein.h)
<i>Arabidopsis</i> callus	Col-0 wild-type	0.037
	CKX4-37	0.244
	CKX4-40	0.258
	CKX4-41	0.320

10

3. Plant phenotypic analysis

The phenotypes generated by overexpression of the *AtCKX4* gene in tobacco and *Arabidopsis* are basically similar as those of *AtCKX1* and *AtCKX2* expressing plants, i.e. enhanced rooting, reduced apical dominance, dwarfing
15 and yellowing of intercostal regions in older leaves of tobacco. An additional phenotype in tobacco are lanceolate leaves (altered length-to-wide ratio).

Example 6. Cloning of the *AtCKX5* gene

The following primers are used to PCR amplify the *AtCKX5* gene from
20 *Arabidopsis thaliana*, accession Columbia:

Sequence of 5' primer: ggggtaccGTGAAATGACGTCAAGCTTTCTT (SEQ ID NO:21)

Sequence of 3' primer: ggggtaccCTTTCCTCTTGGTTTTGTCCTGT (SEQ ID NO:22)

25

A 2833-bp PCR fragment, produced by this PCR amplification, is inserted as a blunt-end product in pCR-Blunt II-TOPO cloning vector (Invitrogen).

Example 7. Cloning of the AtCKX6 gene

- 5 The following primers are used to PCR amplify the AtCKX6 gene from *Arabidopsis thaliana*, accession Columbia:

Sequence of 5' primer: gctctagaTCAGGAAAAGAACCATGCTTATAG (SEQ ID NO:23)

- 10 Sequence of 3' primer: gctctagaTCATGAGTATGAGACTGCCTTTTG (SEQ ID NO:24)

A 1949-bp PCR fragment, produced by this PCR amplification, is inserted as a blunt-end product in pCR-Blunt II-TOPO cloning vector (Invitrogen).

- 15 **Example 8. Expression of the AtCKX2 gene under a root-specific promoter in transgenic tobacco leads to increased root production**

The *AtCKX2* gene (see example 3) is cloned under control of the WRKY6 promoter of *Arabidopsis*, which is a promoter that drives root-specific expression.

- 20 Transgenic plants expressing the *AtCKX2* specifically in the roots show increased root production without negatively affecting shoot development and without inducing premature leaf senescence.

Example 9. Tobacco seedling growth test

- 25 Seeds of *AtCKX1*-, *AtCKX2*-overexpressing and wild-type tobacco were sown in vitro, brought to culture room after 4 days (after cold treatment), and germinated after 6 days.

Observation on seedling growth were made 13 days after sowing:

- 30 Wild type:
just two cotyledons, tiny leaves visible;

root length 9-11 mm, no side root, no adventitious root

AtCKX1:

- clone 50 (strongest phenotype in green house): root length 18-21 mm, 1-2 side
5 branches (sb), 1-2 adventitious roots (ar)
clone 8: root length 13-16 mm, 1 ar, 1 sb
clone 2: root length 17-20 mm, 1-2 ar, 1-2 sb

AtCKX2:

- 10 clone 2: root length 14-18 mm
clone 38: root length 16-19 mm
clone 40: root length 15-20 mm
clone 2 (very low expresser): root length 10 mm
15 formation of more adventitious roots also evident for *AtCKX2* transgenics

***AtCKX1* and *AtCKX2* plants, general observations:**

- longer primary root (up to 100%!) which has already branched (1-2 side
branches), more adventitious roots (1-2)
20 Primary true leaves are advanced, cotyledons are somewhat more expanded.

These results show that overexpression of cytokinin oxidase enhances the
growth and development of both the main root and the adventitious roots,
resulting in early vigor.

25

30

REFERENCES

- Alberts, B., Bray, D., Lewis, J., Raff, M., Roberts, K., and Watson, J. D. (1994). "Molecular Biology of the Cell." Garland Publishing Inc..
- 5 An, G., Watson, B. D., Stachel, S., Gordon, M. P., and Nester, E. W. (1985). New cloning vehicles for transformation of higher plants. *EMBO J.* **4**, 277-284.
- Armstrong, C. L., Petersen, W. P., Buchholz, W. G., Bowen, B. A., and Sulc, S. L. (1990). Factors affecting PEG-mediated stable transformation of maize protoplasts. *Plant Cell Reports* **9**, 335-339.
- 10 Banerjee, A., Pramanik, A., Bhattacharjya, S., and Balaram, P. (1996). Omega amino acids in peptide design: incorporation into helices. *Biopolymers* **39**, 769-777.
- Baron, M. H. and Baltimore, D. (1982). Antibodies against the chemically synthesized genome-linked protein of poliovirus react with native virus-specific proteins. *Cell* **28**, 395-404.
- 15 Bartel, P. L. and Fields, S. (1997). "The Yeast Two-Hybrid System." Oxford University Press.
- Benkirane, N., Guichard, G., Briand, J. P., and Muller, S. (1996). Exploration of requirements for peptidomimetic immune recognition. Antigenic and immunogenic properties of reduced peptide bond pseudopeptide analogues of a histone hexapeptide. *J. Biol Chem.* **271**, 33218-33224.
- 20 Berry, A. and Brenner, S. E. (1994). A prototype computer system for de novo protein design. *Biochem. Soc. Trans.* **22**, 1033-1036.
- Christou, P., McCabe, D. E., and Swain, W. F. (1988). Stable transformation of soybean callus by DNA-coated gold particles. *Plant Physiol.* **87**, 671-674.
- 25 Crossway, A., Oakes, J. V., Irvine, J. M., Ward, B., Knauf, V. C., and Shewmaker, C. K. (1986). Integration of foreign DNA following microinjection of tobacco mesophyll protoplasts. *Mol. Gen. Genet.* **202**, 179-185.

- Dale, E. C. and Ow, D. W. (1990). Intra- and intermolecular site-specific recombination in plant cells mediated by bacteriophage P1 recombinase. *Gene* **91**, 79-85.
- Dodds, J. H. (1985). "Plant genetic engineering." Cambridge University Press.
- 5 Doerner, P., Jorgensen, J. E., You, R., Steppuhn, J., and Lamb, C. (1996). Control of root growth and development by cyclin expression. *Nature* **380**, 520-523.
- Dorner, B., Husar, G. M., Ostresh, J. M., and Houghten, R. A. (1996). The synthesis of peptidomimetic combinatorial libraries through successive amide
10 alkylations. *Bioorg.Med.Chem.* **4**, 709-715.
- Ellis, J. G., Llewellyn, D. J., Dennis, E. S., and Peacock, W. J. (1987). Maize Adh-1 promoter sequences control anaerobic regulation: addition of upstream promoter elements from constitutive genes is necessary for expression in tobacco. *EMBO J.* **6**, 11-16.
- 15 Fassina, G. and Melli, M. (1994). Identification of interactive sites of proteins and protein receptors by computer-assisted searches for complementary peptide sequences. *Immunomethods.* **5**, 114-120.
- Fedoroff, N. V. and Smith, D. L. (1993). A versatile system for detecting transposition in Arabidopsis. *Plant J.* **3**, 273-289.
- 20 Hanahan, D. (1983). Studies on transformation of Escherichia coli with plasmids. *J.Mol.Biol* **166**, 557-580.
- Hansen, G. and Chilton, M. D. (1996). "Agrolistic" transformation of plant cells: integration of T-strands generated in planta. *Proc.Natl.Acad.Sci.U.S.A* **93**, 14978-14983.
- 25 Hansen, G., Shillito, R. D., and Chilton, M. D. (1997). T-strand integration in maize protoplasts after codelivery of a T-DNA substrate and virulence genes. *Proc.Natl.Acad.Sci.U.S.A* **94**, 11726-11730.

- Hanson, B., Engler, D., Moy, Y., Newman, B., Ralston, E., and Gutterson, N. (1999). A simple method to enrich an *Agrobacterium*-transformed population for plants containing only T-DNA sequences. *Plant J.* **19**, 727-734.
- Harlow, E. and Lane, D. (1988). "Antibodies: A Laboratory Manual." Cold Spring Harbor Laboratory Press.
- Herrera-Estrella, L., De Block, M., Messens, E. H. J. P., Van Montagu, M., and Schell, J. (1983). Chimeric genes as dominant selectable markers in plant cells. *EMBO J.* **2**, 987-995.
- Hoffman, D. L., Laiter, S., Singh, R. K., Vaisman, I. I., and Tropsha, A. (1995). Rapid protein structure classification using one-dimensional structure profiles on the bioSCAN parallel computer. *Comput.Appl.Biosci.* **11**, 675-679.
- Krens, F. A., Molendijk, L., Wullems, G. J., and Schilperoort, R. A. (1982). *In vitro* transformation of plant protoplasts with Ti-plasmid DNA. *Nature* **296**, 72-74.
- Lerner, R. A. (1982). Tapping the immunological repertoire to produce antibodies of predetermined specificity. *Nature* **299**, 593-596.
- Lerner, R. A., Green, N., Alexander, H., Liu, F. T., Sutcliffe, J. G., and Shinnick, T. M. (1981). Chemically synthesized peptides predicted from the nucleotide sequence of the hepatitis B virus genome elicit antibodies reactive with the native envelope protein of Dane particles. *Proc.Natl.Acad.Sci.U.S.A* **78**, 3403-3407.
- Liddle, J. E. and Cryer, A. (1991). "A Practical Guide to Monoclonal Antibodies." Wiley New York.
- Loffler, J., Langui, D., Probst, A., and Huber, G. (1994). Accumulation of a 50 kDa N-terminal fragment of beta-APP695 in Alzheimer's disease hippocampus and neocortex. *Neurochem.Int.* **24**, 281-288.
- Monge, A., Lathrop, E. J., Gunn, J. R., Shenkin, P. S., and Friesner, R. A. (1995). Computer modeling of protein folding: conformational and energetic analysis of reduced and detailed protein models. *J.Mol.Biol* **247**, 995-1012.

- Murakami, T., Simonds, W. F., and Spiegel, A. M. (1992). Site-specific antibodies directed against G protein beta and gamma subunits: effects on alpha and beta gamma subunit interaction. *Biochemistry* **31**, 2905-2911.
- Olszewski, K. A., Kolinski, A., and Skolnick, J. (1996). Folding simulations and
5 computer redesign of protein A three-helix bundle motifs. *Proteins* **25**, 286-299.
- Osborne, B. I., Wirtz, U., and Baker, B. (1995). A system for insertional mutagenesis and chromosomal rearrangement using the Ds transposon and Cre-lox. *Plant J.* **7**, 687-701.
- Ostresh, J. M., Blondelle, S. E., Dorner, B., and Houghten, R. A. (1996).
10 Generation and use of nonsupport-bound peptide and peptidomimetic combinatorial libraries. *Methods Enzymol.* **267**, 220-234.
- Pabo, C. O. and Suchanek, E. G. (1986). Computer-aided model-building strategies for protein design. *Biochemistry* **25**, 5987-5991.
- Paszkowski, J., Shillito, R. D., Saul, M., Mandak, V., and Hohn, T. H. B. P. I.
15 (1984). Direct gene transfer to plants. *EMBO J.* **3**, 2717-2722.
- Peralta, E. G., Hellmiss, R., and Ream, W. (1986). Overdrive, a T-DNA transmission enhancer on the *A. tumefaciens* tumour-inducing plasmid. *EMBO J.* **5**, 1137-1142.
- Renouf, D. V. and Hounsell, E. F. (1995). Molecular modelling of glycoproteins
20 by homology with non-glycosylated protein domains, computer simulated glycosylation and molecular dynamics. *Adv.Exp.Med.Biol* **376**, 37-45.
- Rose, R. B., Craik, C. S., Douglas, N. L., and Stroud, R. M. (1996). Three-dimensional structures of HIV-1 and SIV protease product complexes. *Biochemistry* **35**, 12933-12944.
- Rutenber, E. E., McPhee, F., Kaplan, A. P., Gallion, S. L., Hogan, J. C., Jr.,
25 Craik, C. S., and Stroud, R. M. (1996). A new class of HIV-1 protease inhibitor: the crystallographic structure, inhibition and chemical synthesis of an aminimide peptide isostere. *Bioorg.Med.Chem.* **4**, 1545-1558.

- Sambrook, J., Fritsch, E. F., and Maniatis, T. (1989). "Molecular Cloning: A Laboratory Manual." Cold Spring Harbor Laboratory Press.
- Schlappi, M., Smith, D., and Fedoroff, N. (1993). TnpA trans-activates methylated maize Suppressor-mutator transposable elements in transgenic tobacco. *Genetics* **133**, 1009-1021.
- Shioda, T., Andriole, S., Yahata, T., and Isselbacher, K. J. (2000). A green fluorescent protein-reporter mammalian two-hybrid system with extrachromosomal maintenance of a prey expression plasmid: Application to interaction screening. *Proc.Natl.Acad.Sci.U.S.A* **97**, 5220-5224.
- 10 Tamura, R. N., Cooper, H. M., Collo, G., and Quaranta, V. (1991). Cell type-specific integrin variants with alternative alpha chain cytoplasmic domains. *Proc.Natl.Acad.Sci.U.S.A* **88**, 10183-10187.
- Van Haaren, M. J., Sedee, N. J., Schilperoort, R. A., and Hooykaas, P. J. (1987). Overdrive is a T-region transfer enhancer which stimulates T-strand production in *Agrobacterium tumefaciens*. *Nucleic Acids Res.* **15**, 8983-8997.
- 15 Van Sluys, M. A., Tempe, J., and Fedoroff, N. (1987). Studies on the introduction and mobility of the maize Activator element in *Arabidopsis thaliana* and *Daucus carota*. *EMBO J.* **6**, 3881-3889.
- Wang, K., Genetello, C., Van Montagu, M., and Zambryski, P. C. (1987). Sequence context of the T-DNA border repeat element determines its relative activity during T-DNA transfer to plant cells. *Mol.Gen.Genet.* **210**, 338-346.
- 20 Woulfe, J., Lafortune, L., de Nadai, F., Kitabgi, P., and Beaudet, A. (1994). Post-translational processing of the neurotensin/neuromedin N precursor in the central nervous system of the rat—II. Immunohistochemical localization of maturation products. *Neuroscience* **60**, 167-181.
- 25 Zhang, Y. L., Dawe, A. L., Jiang, Y., Becker, J. M., and Naider, F. (1996). A superactive peptidomimetic analog of a farnesylated dodecapeptide yeast pheromone. *Biochem.Biophys.Res.Comm.* **224**, 327-331.

THIS PAGE BLANK (USPTO)

CLAIMS

1. A method for stimulating root growth and/or enhancing the formation of lateral and adventitious roots and/or altering root geotropism comprising expression
5 of a plant cytokinin oxidase or comprising expression of another protein that reduces the level of active cytokinins in plants or plant parts.
2. An isolated nucleic acid encoding a novel plant protein having cytokinin oxidase activity or encoding an immunologically active and/or functional fragment of such a protein selected from the group consisting of:
 - 10 (a) nucleic acid sequences comprising a DNA sequence as given in any of SEQ ID NOs 1, 3, 5, 7, 9 or 11, or the complement thereof,
 - (b) nucleic acid sequences comprising the RNA sequences corresponding to any of SEQ ID NOs 1, 3, 5, 7, 9 or 11, or the complement thereof,
 - (c) nucleic acid sequences hybridizing to the nucleotide sequence as defined
15 in (a) or (b),
 - (d) nucleic acid sequences encoding a protein with an amino acid sequence which is at least 35% similar to the amino acid sequence as given in SEQ ID NO 2,
 - (e) nucleic acid sequences encoding a protein with an amino acid sequence
20 which is at least 40% similar to the amino acid sequence as given in SEQ ID NO 4,
 - (f) nucleic acid sequences encoding a protein with an amino acid sequence which is at least 42% similar to the amino acid sequence as given in SEQ ID NO 6,
 - 25 (g) nucleic acid sequences encoding a protein with an amino acid sequence which is at least 44% similar to the amino acid sequence as given in SEQ ID NO 8,
 - (h) nucleic acid sequences encoding a protein with an amino acid sequence
30 which is at least 40% similar to the amino acid sequence as given in SEQ ID NO 10,

- (i) nucleic acid sequences encoding a protein with an amino acid sequence which is at least 40% similar to the amino acid sequence as given in SEQ ID NO 12,
 - (j) nucleic acid sequences encoding a protein comprising the amino acid sequence as given in any of SEQ ID NOs 2, 4, 6, 8, 10 or 12,
 - (k) nucleic acid sequences which are degenerated as a result of the genetic code to a nucleotide sequence of a nucleic acid as given in any of SEQ ID NOs 1, 3, 5, 7, 9 or 11, or as defined in (a) to (j),
 - (l) nucleic acid sequences which are diverging due to the differences in codon usage between the organisms to a nucleotide sequence encoding a protein as given in any of SEQ ID NOs 2, 4, 6, 8, 10, 12 or as defined in (a) to (j),
 - (m) nucleic acid sequences which are diverging due to the differences between alleles encoding a protein as given in SEQ ID NOs 2, 4, 6, 8, 10 or 12, or as defined in (a) to (j),
 - (n) nucleic acid sequences encoding an immunologically active and/or functional fragment of a protein encoded by a DNA sequence as given in any of SEQ ID NOs 1, 3, 5, 7, 9 or 11, or a functional fragment of a nucleic acid as defined in any one of (a) to (m), and,
 - (o) nucleic acid sequences encoding a protein as defined in SEQ ID NO 2, 4, 6, 8, 10 or 12 or a nucleic acid as defined in any one of (a) to (n) characterized in that said sequence is a cDNA sequence.
3. An isolated nucleic acid according to claim 2 which is DNA, cDNA, genomic DNA or synthetic DNA, or RNA wherein T is replaced by U.
 4. A nucleic acid molecule of at least 15 nucleotides in length hybridizing specifically with a nucleic acid of claim 2 or 3.
 5. A nucleic acid molecule of at least 15 nucleotides in length specifically amplifying a nucleic acid of claim 2 or 3.
 6. A vector comprising a nucleic acid sequence according to claim 2 or 3.

7. A vector according to claim 6 which is an expression vector wherein the nucleic acid sequence is operably linked to one or more control sequences allowing the expression of said sequence in prokaryotic and/or eukaryotic host cells.
- 5 8. A host cell containing a nucleic acid molecule according to claim 2 or 3 or a vector according to claim 6 or 7.
9. The host cell of claim 8, wherein the host cell is a bacterial, insect, fungal, plant or animal cell.
10. An isolated polypeptide encodable by a nucleic acid of claim 2 or 3, or a
10 homologue or a derivative thereof, or an immunologically active and/or functional fragment thereof.
11. The polypeptide of claim 10 which has an amino acid sequence as given in SEQ ID NO 2, 4, 6, 8, 10 or 12, or a homologue or a derivative thereof, or an immunologically active and/or functional fragment thereof.
- 15 12. A method for producing a polypeptide according to claim 10 or 11 comprising culturing a host cell of claim 8 or 9 under conditions allowing the expression of the polypeptide and recovering the produced polypeptide from the culture.
13. An antibody specifically recognizing a polypeptide of claim 10 or 11 or a specific epitope thereof.
- 20 14. A method for the production of transgenic plants, plant cells or plant tissues comprising the introduction of a nucleic acid molecule according to any of claims 2 or 3 in an expressible format or a vector according to claim 6 or 7 in said plant, plant cell or plant tissue.
- 25 15. A method for the production of altered plants, plant cells or plant tissues comprising the introduction of a polypeptide of claim 10 or 11 directly into a cell, a tissue or an organ of said plant.
16. A method for effecting the expression of a polypeptide of claim 10 or 11 comprising the introduction of a nucleic acid molecule of claim 2 or 3 operably

linked to one or more control sequences or a vector according to claim 6 or 7 stably into the genome of a plant cell.

17. The method of claim 15 or 16 further comprising regenerating a plant from said plant cell.

5 18. A transgenic plant cell comprising a nucleic acid sequence of claim 2 or 3 which is operably linked to regulatory elements allowing transcription and/or expression of said nucleic acid in plant cells or obtainable by a method of claim 15 or 16.

10 19. The transgenic plant cell of claim 18 wherein said nucleic acid of claim 2 or 3 is stably integrated into the genome of said plant cell.

20. A transgenic plant or plant tissue comprising plant cells of claim 18 or 19.

21. A harvestable part of a plant of claim 20.

22. The harvestable part of claim 21 which is selected from the group consisting of seeds, leaves, fruits, stem cultures, rhizomes and bulbs.

15 23. The progeny derived from any of the plants or plant parts of any of claims 20 to 22.

24. A method for stimulating root growth comprising expression of a nucleic acid of claim 2 or 3 or comprising expression of another protein that reduces the level of active cytokinins in plants or plant parts.

20 25. A method for enhancing the formation of lateral and adventitious roots comprising expression of a nucleic acid of claim 2 or 3 or comprising expression of another protein that reduces the level of active cytokinins in plants or plant parts.

25 26. A method for altering root geotropism comprising altering the expression of a nucleic acid of claim 2 or 3 or comprising expression of another protein that reduces the level of active cytokinins in plants or plant parts.

27. A method of any of claims 24 to 26, said method leading to an increase in yield.

28. The method of any of claims 24 to 27 wherein said expression of said nucleic acid occurs under the control of a strong constitutive promoter.
29. The method of any of claims 24 to 27 wherein said expression of said nucleic acid occurs under the control of a promoter that is preferentially expressed in roots.
30. A method for identifying and obtaining proteins interacting with a polypeptide of claim 10 or 11 comprising a screening assay wherein a polypeptide of claim 10 or 11 is used.
31. The method of claim 30 comprising a two-hybrid screening assay wherein a polypeptide of claim 10 or 11 as a bait and a cDNA library as prey are used.
32. A method for modulating the interaction between a polypeptide of claim 10 or 11 and interacting protein partners obtainable by a method according to claim 30 or 31.
33. A method for identifying and obtaining compounds interacting with a polypeptide of claim 10 or 11 comprising the steps of:
- (a) providing a two-hybrid system wherein a polypeptide of claim 10 or 11 and an interacting protein partner obtainable by a method according to claim 30 or 31 are expressed,
 - (b) interacting said compound with the complex formed by the expressed polypeptides as defined in a), and,
 - (c) performing measurement of interaction of said compound with said polypeptide or the complex formed by the expressed polypeptides as defined in (a).
34. A method for identifying compounds or mixtures of compounds which specifically bind to a polypeptide of claim 10 or 11, comprising:
- (a) combining a polypeptide of claim 10 or 11 with said compound or mixtures of compounds under conditions suitable to allow complex formation, and,
 - (b) detecting complex formation, wherein the presence of a complex identifies a compound or mixture which specifically binds said polypeptide.

35. A method of any of claims 30 to 34 wherein said compound or mixture inhibits the activity of said polypeptide of claim 10 or 11 and can be used for the rational design of chemicals.
36. Use of a compound or mixture identified by means of a method of any of claims 30 to 34 as a plant growth regulator or herbicide.
37. A method for production of a plant growth regulator or herbicide composition comprising the steps of the method of any of claims 30 to 34 and formulating the compounds obtained from said steps in a suitable form for the application in agriculture or plant cell or tissue culture.
38. A method for the design of or screening for growth-promoting chemicals or herbicides comprising the use of a nucleic acid of claim 2 or 3 or a vector of claim 6 or 7.
39. Use of a nucleic acid molecule of claim 2 or 3, the vector of claim 6 or 7, a polypeptide of claim 10 or 11 for increasing yield.
40. Use of a nucleic acid molecule of claim 2 or 3, the vector of claim 6 or 7, a polypeptide of claim 10 or 11 for stimulating root growth.
41. Use of a nucleic acid molecule of claim 2 or 3, the vector of claim 6 or 7, a polypeptide of claim 10 or 11 for enhancing the formation of lateral and adventitious roots.
42. Use of a nucleic acid molecule of claim 2 or 3, the vector of claim 6 or 7, a polypeptide of claim 10 or 11 for altering root geotropism.
43. Diagnostic composition comprising at least a nucleic acid molecule of any of claims 2 to 5, the vector of claim 6 or 7, a polypeptide of claim 10 or 11 or an antibody of claim 13.

ABSTRACT

The present invention relates to methods for stimulating root growth and/or enhancing the formation of lateral and adventitious roots and/or altering root geotropism comprising expression of a plant cytokinin oxidase or comprising
5 expression of another protein that reduces the level of active cytokinins in plants or plant parts.

The invention also relates to novel plant cytokinin oxidase proteins, nucleic acid sequences encoding cytokinin oxidase proteins as well as to vectors, host cells,
10 transgenic cells and plants comprising said sequences. The invention also relates to the use of said sequences for increasing yield and/or enhancing early vigor and/or modifying root/shoot ratio and/or improving lodging and/or increasing drought tolerance and/or promoting in vitro propagation of explants and/or modifying cell fate and/or plant development and/or plant morphology and/or
15 plant biochemistry and/or plant physiology. The invention also relates to the use of said sequences in the above mentioned methods.

The invention also relates to methods for identifying and obtaining proteins and compounds interacting with cytokinin oxidase proteins. The invention also relates to the use of said compounds as a plant growth regulator or herbicide.

THIS PAGE BLANK (USPTO)




FIGURE 1

2/8

ZmCKX1	1	1	5	10	15	20	25	30	35	40	45	50	55	60	65	70	75	80	85	90	95	100
AtCKX1	1	1	5	10	15	20	25	30	35	40	45	50	55	60	65	70	75	80	85	90	95	100
AtCKX2	1	1	5	10	15	20	25	30	35	40	45	50	55	60	65	70	75	80	85	90	95	100
AtCKX3	1	1	5	10	15	20	25	30	35	40	45	50	55	60	65	70	75	80	85	90	95	100
AtCKX4	1	1	5	10	15	20	25	30	35	40	45	50	55	60	65	70	75	80	85	90	95	100
ZmCKX1	82	101	110	120	130	140	150	160	170	180	190	200										
AtCKX1	101	110	120	130	140	150	160	170	180	190	200											
AtCKX2	70	110	120	130	140	150	160	170	180	190	200											
AtCKX3	83	110	120	130	140	150	160	170	180	190	200											
AtCKX4	77	110	120	130	140	150	160	170	180	190	200											
ZmCKX1	173	201	210	220	230	240	250	260	270	280	290	300										
AtCKX1	190	210	220	230	240	250	260	270	280	290	300											
AtCKX2	154	210	220	230	240	250	260	270	280	290	300											
AtCKX3	171	210	220	230	240	250	260	270	280	290	300											
AtCKX4	177	210	220	230	240	250	260	270	280	290	300											
ZmCKX1	273	301	310	320	330	340	350	360	370	380	390	400										
AtCKX1	290	310	320	330	340	350	360	370	380	390	400											
AtCKX2	254	310	320	330	340	350	360	370	380	390	400											
AtCKX3	271	310	320	330	340	350	360	370	380	390	400											
AtCKX4	277	310	320	330	340	350	360	370	380	390	400											
ZmCKX1	373	401	410	420	430	440	450	460	470	480	490	500										
AtCKX1	377	410	420	430	440	450	460	470	480	490	500											
AtCKX2	340	410	420	430	440	450	460	470	480	490	500											
AtCKX3	358	410	420	430	440	450	460	470	480	490	500											
AtCKX4	363	410	420	430	440	450	460	470	480	490	500											
ZmCKX1	466	501	510	520	530	540	550	560	570	580	590	600										
AtCKX1	473	510	520	530	540	550	560	570	580	590	600											
AtCKX2	435	510	520	530	540	550	560	570	580	590	600											
AtCKX3	451	510	520	530	540	550	560	570	580	590	600											
AtCKX4	457	510	520	530	540	550	560	570	580	590	600											

FIGURE 2






FIGURE 3

BEST AVAILABLE COPY




FIGURE 4




FIGURE 5

BEST AVAILABLE COPY

BEST AVAILABLE COPY




FIGURE 6




FIGURE 7

BEST AVAILABLE COPY

BEST AVAILABLE COPY



FIGURE 8

SEQUENCE LISTING

<110> CropDesign N.V.

<120> Method for modifying plant morphology, biochemistry and physiology

<130> CROP-005-EP

<140>

<141>

<160> 24

<170> PatentIn Ver. 2.1

<210> 1

<211> 2236

<212> DNA

<213> Arabidopsis thaliana

<400> 1

```

atgggattga cctcatcctt acggttccat agacaaaaca acaagacttt cctcggaaac 60
ttcatgatct tagttctaag ctgtatacca ggtagaacca atctttgttc caatcattct 120
gtagtagacc caaaagaatt accttcttca aatccttcag atattcggtc ctcattagtt 180
tacttagatt tggaggggta tataagcttc gacgatgtcc acaatgtggc caaggacttt 240
ggcaacagat accagttacc acctttggca attctacatc caaggtcagt ttttgatatt 300
tcatcgatga tgaagcatat agtacatctg ggctccacct caaatcttac agtagcagct 360
agaggccatg gtcactcgct tcaaggacaa gctctagctc atcaagggtg tgtcatcaaa 420
atggagtcac ttcgaagtc tgaatcagg atttataagg ggaagcaacc atatgttgat 480
gtctcagggt gtgaaatatg gataaacatt ctacgcgaga ctctaaaata cggctcttca 540
ccaaagtcct ggacagacta ccttcatttg accgttggag gtacactatc taatgctgga 600
atcagcggtc aagcattcaa gcatggaccc caaatcaaca acgtctacca gctagagatt 660
gttacaggta tttcattcat gctttatctc tgcggtagtc tcaaaaaaat atgcacctgt 720
aaagaatatc catctcttca tgagcaaaaa cactgacgac tttaaataat ttttgactat 780
aaaacaagag tgcataggca caaatgtgaa atatgcaaca cacaattgta acttgcacca 840
agaaaaaagt tataaaaaa aacaactgat aagcaatata tttccaatat ttaatcaggg 900
aaaggagaag tcgtaacctg ttctgagaag cgggaattctg aacttttctt cagtgttctt 960
ggcgggcttg gacagtttgg cataatcacc cgggcacgga tctctcttga accagcaccg 1020
catatggtaa agttctatct tgaacaaagt tcaacaata tacgctatga ttctaagaac 1080
cactttcctg acacagtcaa ataactttta atagggttaa tggatcaggg tactctactc 1140
tgacttttct gcattttcaa gggaccaaga atatctgatt tcgaaggaga aaacttttga 1200
ttacgttgaa ggatttgtga taatcaatag aacagacctt ctcaataatt ggcgatcgct 1260
attcagtcct aacgattcca cacaggcaag cagattcaag tcagatggga aaactcttta 1320
ttgcctagaa gtggtcaaat atttcaacct agaagaagct agctctatgg atcaggtaag 1380
atgtgaaagc aatatataac tagacttagt ttccacagag agctccaaat caaccgttgg 1440
ctactagcct actaacataa tgaatgggtg ccgtgcagga aactggcaag ttactttcag 1500
agttaatta tattccatcc actttgtttt catctgaagt gccatatatc gagtttctgg 1560

```

atcgcgtgca tatcgcagag agaaaactaa gagcaaaggg tttatgggag gttccacatc 1620
 cctggctgaa tctcctgatt cctaagagca gcatatacca atttgctaca gaagttttca 1680
 acaacattct cacaagcaac aacaacggtc ctatccttat ttatccagtc aatcaatcca 1740
 agtaagttag caaaatgccca aaagcaaagt cgtccagtgga ttctgaaaca taaattacta 1800
 accatatcca acattttgtg gtttcaggtg gaagaaacat acatctttga taactccaaa 1860
 tgaagatata ttctatctcg tagcctttct cccctctgca gtgccaaatt cctcagggaa 1920
 aaacgatcta gagtaccttt tgaaacaaaa ccaaagagtt atgaacttct gcgcagcagc 1980
 aaacctcaac gtgaagcagt atttgcccca ttatgaaact caaaaagagt ggaaatcaca 2040
 ctttggcaaaa agatgggaaa catttgccaca gaggaaacaa gcctacgacc ctctagcgat 2100
 tctagcacct ggccaaagaa tattccaaaa gacaacagga aaattatctc ccatccaact 2160
 cgcaaagtca aaggcaacag gaagtctca aaggtaccat tacgcatcaa tactgccgaa 2220
 acctagaact gtataa 2236

<210> 2

<211> 575

<212> PRT

<213> Arabidopsis thaliana

<400> 2

Met Gly Leu Thr Ser Ser Leu Arg Phe His Arg Gln Asn Asn Lys Thr
 1 5 10 15

Phe Leu Gly Ile Phe Met Ile Leu Val Leu Ser Cys Ile Pro Gly Arg
 20 25 30

Thr Asn Leu Cys Ser Asn His Ser Val Ser Thr Pro Lys Glu Leu Pro
 35 40 45

Ser Ser Asn Pro Ser Asp Ile Arg Ser Ser Leu Val Ser Leu Asp Leu
 50 55 60

Glu Gly Tyr Ile Ser Phe Asp Asp Val His Asn Val Ala Lys Asp Phe
 65 70 75 80

Gly Asn Arg Tyr Gln Leu Pro Pro Leu Ala Ile Leu His Pro Arg Ser
 85 90 95

Val Phe Asp Ile Ser Ser Met Met Lys His Ile Val His Leu Gly Ser
 100 105 110

Thr Ser Asn Leu Thr Val Ala Ala Arg Gly His Gly His Ser Leu Gln
 115 120 125

Gly Gln Ala Leu Ala His Gln Gly Val Val Ile Lys Met Glu Ser Leu
 130 135 140

Arg Ser Pro Asp Ile Arg Ile Tyr Lys Gly Lys Gln Pro Tyr Val Asp

145	150	155	160
Val Ser Gly Gly Glu Ile Trp Ile Asn Ile Leu Arg Glu Thr Leu Lys	165	170	175
Tyr Gly Leu Ser Pro Lys Ser Trp Thr Asp Tyr Leu His Leu Thr Val	180	185	190
Gly Gly Thr Leu Ser Asn Ala Gly Ile Ser Gly Gln Ala Phe Lys His	195	200	205
Gly Pro Gln Ile Asn Asn Val Tyr Gln Leu Glu Ile Val Thr Gly Lys	210	215	220
Gly Glu Val Val Thr Cys Ser Glu Lys Arg Asn Ser Glu Leu Phe Phe	225	230	235
Ser Val Leu Gly Gly Leu Gly Gln Phe Gly Ile Ile Thr Arg Ala Arg	245	250	255
Ile Ser Leu Glu Pro Ala Pro His Met Val Lys Trp Ile Arg Val Leu	260	265	270
Tyr Ser Asp Phe Ser Ala Phe Ser Arg Asp Gln Glu Tyr Leu Ile Ser	275	280	285
Lys Glu Lys Thr Phe Asp Tyr Val Glu Gly Phe Val Ile Ile Asn Arg	290	295	300
Thr Asp Leu Leu Asn Asn Trp Arg Ser Ser Phe Ser Pro Asn Asp Ser	305	310	315
Thr Gln Ala Ser Arg Phe Lys Ser Asp Gly Lys Thr Leu Tyr Cys Leu	325	330	335
Glu Val Val Lys Tyr Phe Asn Pro Glu Glu Ala Ser Ser Met Asp Gln	340	345	350
Glu Thr Gly Lys Leu Leu Ser Glu Leu Asn Tyr Ile Pro Ser Thr Leu	355	360	365
Phe Ser Ser Glu Val Pro Tyr Ile Glu Phe Leu Asp Arg Val His Ile	370	375	380
Ala Glu Arg Lys Leu Arg Ala Lys Gly Leu Trp Glu Val Pro His Pro	385	390	395
Trp Leu Asn Leu Leu Ile Pro Lys Ser Ser Ile Tyr Gln Phe Ala Thr			400

	405		410		415
Glu Val Phe Asn Asn Ile Leu Thr Ser Asn Asn Asn Gly Pro Ile Leu					
	420		425		430
Ile Tyr Pro Val Asn Gln Ser Lys Trp Lys Lys His Thr Ser Leu Ile					
	435		440		445
Thr Pro Asn Glu Asp Ile Phe Tyr Leu Val Ala Phe Leu Pro Ser Ala					
	450		455		460
Val Pro Asn Ser Ser Gly Lys Asn Asp Leu Glu Tyr Leu Leu Lys Gln					
465		470		475	480
Asn Gln Arg Val Met Asn Phe Cys Ala Ala Ala Asn Leu Asn Val Lys					
	485		490		495
Gln Tyr Leu Pro His Tyr Glu Thr Gln Lys Glu Trp Lys Ser His Phe					
	500		505		510
Gly Lys Arg Trp Glu Thr Phe Ala Gln Arg Lys Gln Ala Tyr Asp Pro					
	515		520		525
Leu Ala Ile Leu Ala Pro Gly Gln Arg Ile Phe Gln Lys Thr Thr Gly					
	530		535		540
Lys Leu Ser Pro Ile Gln Leu Ala Lys Ser Lys Ala Thr Gly Ser Pro					
545		550		555	560
Gln Arg Tyr His Tyr Ala Ser Ile Leu Pro Lys Pro Arg Thr Val					
	565		570		575

<210> 3

<211> 2991

<212> DNA

<213> Arabidopsis thaliana

<400> 3

```

atggctaatac ttcgtttaata gatcacttta atcacggttt taatgatcac caaatcatca 60
aacggtatta aaattgattt acctaaatcc cttaacctca cctctcttac cgatccttcc 120
atcatctccg cagcctctca tgacttcgga aacataacca ccgtgacccc cggcggcgta 180
atctgcccct cctccaccgc tgatatctct cgtctcctcc aatacgccgc aaacggaaaa 240
agtacattcc aagtagcggc tcgtggccaa ggccactcct taaacggcca agcctcggtc 300
tccggcggag taatcgtcaa catgacgtgt atcactgacg tgggtggtttc aaaagacaag 360
aagtagctg acgtggcggc cgggacgtta tgggtggatg tgcttaagaa gacggcggag 420
aaaggggtgt cgccgggttc ttggacggat tatttgcata taaccgtcgg aggaacgttg 480

```

tcgaatggtg gaattggtgg tcaagtgttt cgaaacggtc ctcttggttag taacgtcctt 540
 gaattggacg ttattactgg tacgcatctt ctaaactttg atgtacatac aacaacaaaa 600
 actgtttttg ttttatagta ttttcatatt tttgtaccat aggttttatg ttttatagtt 660
 gtgctaaact tcttgacca cacgtaagtc ttcgaaacac aaaatgcgta acgcatctat 720
 atgttttttg tacatattga atgttggtca tgagaaataa agtaattaca tatacacaca 780
 tttattgtcg tacatatata aataattaaa gacaaatttt cacaattggg agcgtgttaa 840
 tttgggattt ttgtaatgta catgcatgac gcatgcatat ggagcttttc ggttttctta 900
 gatttgtgta gtatttcaaa tatatcattt attttcttct gaataaagag gtggtatatt 960
 tttaaaatag caacatttca gaatttttct ttgaatttac actttttaa ttgttattgt 1020
 taatatggat tttgaataaa taatttcagg gaaagggtgaa atgttgacat gctcgcgaca 1080
 gctaaaccca gaattgttct atggagtggt aggaggttg ggtcaatttg gaattataac 1140
 gagagccaga attgttttgg accatgcacc taaacgggta cgtatcatca tattttacca 1200
 tttgttttag tcagcattca tttttcatta gtaattccgt ttcaatttct aaattttttt 1260
 agtcaataga aaatgattct tatgtcagag cttgattatt tagtgatttt tattgagata 1320
 aaataaaaata taacctaacg gaaataatta ttttactaat cggataatgt ctgattaaaa 1380
 cattttatga tattacacta agagagtttag agacgtatgg atcacaaaac atgaagcttt 1440
 cttagatggg atcctaaaaac taaagttagg tacaagtttg gaatttaggt caaatgctta 1500
 agttgcatta atttgaacaa aatctatgca ttgaataaaa aaaagatatg gattatttta 1560
 taaagtatag tccttgtaat cctaggactt gttgtcta atctgtcttat gcgtgcaaat 1620
 ctttttgatg tcaatatata atccttgttt attagagtca agctctttca ttagtcaact 1680
 actcaaatat actccaaagt ttagaatata gtcttctgac taattagaat cttacaaccg 1740
 ataaacgtta caatttgggt atcattttta aaaacagatt tggtcataat atacgatgac 1800
 gttctgtttt agtttcatct attcacaat tttatataat tattttcaag aaaatattga 1860
 aatactatac tgtaatatgg tttctttata tatgtgtgta taaattaaat gggattgttt 1920
 tctctaaatg aaattgtgta ggccaaatgg tttcggatgc tctacagtga tttcacaact 1980
 tttacaaagg accaagaacg tttgatatca atggcaaacg atattggagt cgactattta 2040
 gaagggtcaa ttttctatc aaacgggtgc gttgacacct cttttttccc accttcagat 2100
 caatctaaag tcgctgatct agtcaagcaa cacggtatca tctatgttct tgaagtagcc 2160
 aagtattatg atgatcccaa tctccccatc atcagcaagg tactacacat ttacattttc 2220
 atcatcgttt ttatcatacc ataagatatt taaatgattc atcattgcac cacattaaga 2280
 tattcatcat catcatcggt acattttttt ttgcatctta tgcttctcat aatctactat 2340
 tgtgtagggt attgacacat taacgaaaac attaatgtac ttgcccgggt tcatatcaat 2400
 gcacgacgtg gcctacttcg atttcttgaa ccgtgtacat gtcgaagaaa ataaactcag 2460
 atctttggga ttatgggaac ttctcatcc ttggcttaac ctctacgttc ctaaactctg 2520
 gattctcgat tttcataacg gtgtgtgcaa agacattctt ctttaagcaa aatcagcttc 2580
 gggactcgct cttctctatc caacaaaccg gaataagtag atacttctct tcattcatat 2640
 ttatcttcaa gaaccaaagt aaataaattt ctatgaactg attatgctgt tattgttaga 2700
 tgggacaatc gtatgtcggc gatgatacca gagatcgatg aagatgttat atatattatc 2760
 ggactactac aatccgctac ccaaaggat cttccagaag tggagagcgt taacgagaag 2820
 ataattaggt tttgcaagga ttcagggtatt aagattaagc aatatcta gcatatact 2880
 agtaaagaag attggattga gcatttttga tcaaaatggg atgatttttc gaagaggaaa 2940
 gatctatttg atcccaagaa actgttatct ccagggcaag acatcttttg a 2991

<210> 4

<211> 501

<212> PRT

<213> Arabidopsis thaliana

<400> 4

Met	Ala	Asn	Leu	Arg	Leu	Met	Ile	Thr	Leu	Ile	Thr	Val	Leu	Met	Ile	
1				5					10					15		
Thr	Lys	Ser	Ser	Asn	Gly	Ile	Lys	Ile	Asp	Leu	Pro	Lys	Ser	Leu	Asn	
			20					25					30			
Leu	Thr	Leu	Ser	Thr	Asp	Pro	Ser	Ile	Ile	Ser	Ala	Ala	Ser	His	Asp	
		35					40					45				
Phe	Gly	Asn	Ile	Thr	Thr	Val	Thr	Pro	Gly	Gly	Val	Ile	Cys	Pro	Ser	
	50						55				60					
Ser	Thr	Ala	Asp	Ile	Ser	Arg	Leu	Leu	Gln	Tyr	Ala	Ala	Asn	Gly	Lys	
65						70				75					80	
Ser	Thr	Phe	Gln	Val	Ala	Ala	Arg	Gly	Gln	Gly	His	Ser	Leu	Asn	Gly	
			85						90					95		
Gln	Ala	Ser	Val	Ser	Gly	Gly	Val	Ile	Val	Asn	Met	Thr	Cys	Ile	Thr	
			100					105					110			
Asp	Val	Val	Val	Ser	Lys	Asp	Lys	Lys	Tyr	Ala	Asp	Val	Ala	Ala	Gly	
	115						120					125				
Thr	Leu	Trp	Val	Asp	Val	Leu	Lys	Lys	Thr	Ala	Glu	Lys	Gly	Val	Ser	
	130					135					140					
Pro	Val	Ser	Trp	Thr	Asp	Tyr	Leu	His	Ile	Thr	Val	Gly	Gly	Thr	Leu	
145					150					155					160	
Ser	Asn	Gly	Gly	Ile	Gly	Gly	Gln	Val	Phe	Arg	Asn	Gly	Pro	Leu	Val	
			165						170					175		
Ser	Asn	Val	Leu	Glu	Leu	Asp	Val	Ile	Thr	Gly	Lys	Gly	Glu	Met	Leu	
		180						185					190			
Thr	Cys	Ser	Arg	Gln	Leu	Asn	Pro	Glu	Leu	Phe	Tyr	Gly	Val	Leu	Gly	
		195					200					205				
Gly	Leu	Gly	Gln	Phe	Gly	Ile	Ile	Thr	Arg	Ala	Arg	Ile	Val	Leu	Asp	
	210					215					220					
His	Ala	Pro	Lys	Arg	Ala	Lys	Trp	Phe	Arg	Met	Leu	Tyr	Ser	Asp	Phe	
225					230					235					240	
Thr	Thr	Phe	Thr	Lys	Asp	Gln	Glu	Arg	Leu	Ile	Ser	Met	Ala	Asn	Asp	

245	250	255
Ile Gly Val Asp Tyr Leu Glu Gly Gln Ile Phe Leu Ser Asn Gly Val 260	265	270
Val Asp Thr Ser Phe Phe Pro Pro Ser Asp Gln Ser Lys Val Ala Asp 275	280	285
Leu Val Lys Gln His Gly Ile Ile Tyr Val Leu Glu Val Ala Lys Tyr 290	295	300
Tyr Asp Asp Pro Asn Leu Pro Ile Ile Ser Lys Val Ile Asp Thr Leu 305	310	315 320
Thr Lys Thr Leu Ser Tyr Leu Pro Gly Phe Ile Ser Met His Asp Val 325	330	335
Ala Tyr Phe Asp Phe Leu Asn Arg Val His Val Glu Glu Asn Lys Leu 340	345	350
Arg Ser Leu Gly Leu Trp Glu Leu Pro His Pro Trp Leu Asn Leu Tyr 355	360	365
Val Pro Lys Ser Arg Ile Leu Asp Phe His Asn Gly Val Val Lys Asp 370	375	380
Ile Leu Leu Lys Gln Lys Ser Ala Ser Gly Leu Ala Leu Leu Tyr Pro 385	390	395 400
Thr Asn Arg Asn Lys Trp Asp Asn Arg Met Ser Ala Met Ile Pro Glu 405	410	415
Ile Asp Glu Asp Val Ile Tyr Ile Ile Gly Leu Leu Gln Ser Ala Thr 420	425	430
Pro Lys Asp Leu Pro Glu Val Glu Ser Val Asn Glu Lys Ile Ile Arg 435	440	445
Phe Cys Lys Asp Ser Gly Ile Lys Ile Lys Gln Tyr Leu Met His Tyr 450	455	460
Thr Ser Lys Glu Asp Trp Ile Glu His Phe Gly Ser Lys Trp Asp Asp 465	470	475 480
Phe Ser Lys Arg Lys Asp Leu Phe Asp Pro Lys Lys Leu Leu Ser Pro 485	490	495
Gly Gln Asp Ile Phe		

<210> 5

<211> 3302

<212> DNA

<213> *Arabidopsis thaliana*

<400> 5

```

atggcgagtt ataatcttcg ttcacaagtt cgtcttatag caataacaat agtaatcatc 60
attactctct caactccgat cacaaccaac acatcaccac aaccatggaa tatectttca 120
cacaacgaat tcgccggaaa actcacctcc tctcctcct ccgtcgaatc agccgccaca 180
gatttcggcc acgtcaccaa aatcttccct tccgccgtct taatcccttc ctccgttgaa 240
gacatcacag atctcataaa actctctttt gactctcaac tgtcttttcc tttagccgct 300
cgtggtcacg gacacagcca ccgtggccaa gcctcggcta aagacggagt tgtggccaac 360
atgcggtcca tggtaaaccg ggatcgaggt atcaagggtg ctaggacctg tttatatgtt 420
gacgtggacg ctgcgtggct atggattgag gtgttgaata aaactttgga gttaggggta 480
acgccggttt cttggacgga ttatttgtat ttaacagtcg gtgggacgtt atcaaacggc 540
ggaattagtg gacaaacgtt tcggtacggg ccacagatca ctaatgttct agagatggat 600
gttattactg gtacgtacca cgatcttttt cacacagaga ttaaaaaaaaa cagtaatagt 660
gatttttaact tcgtacgttt ctgatagaca acaaagaact tcgtacgttt ttcgaagttt 720
tttcgtcttt ttcatttttag atctgcgcgg ccatttttgg ttatgctatt gtttgtttgt 780
attgtttgtc tctgtttatt tatctctcga acttggtgat agcttttctt cttttcacac 840
atcaatctaa tcaccttttt tggctcttaag attagaaaga agatacggac taggtaaaaa 900
taggtgggtg taaacgtaga cgcattaaaa aaatattggt ttttttattt tttgataagc 960
aaaattggtg gttggtctaa gattataaac ttgatattaa tgcaaaggtc gatctagcaa 1020
tagaagatta atcaatattc ttggtgtttt aacaacagat tatttcatca ttaaaatcgt 1080
gaaacaaaga aatttttggt gtatacatta cgtgtagttt tgtagttta ttaaaaaaaaa 1140
tagtatatag ttttggttaa acgcgattta ttagtaaca cattagtata ttacacgttt 1200
aaccaactaa actttttttt ttgaataatt atgttctata tttcttactc aaattatgca 1260
aatttcgtgg attcgaagtc aaatttctgc gaaatttaca tggatcataa ttataaaact 1320
gttcatataa cccggtgaac aaacagacaa ttaagggttt gaatgggtac ggcgggtggg 1380
gcggacacaa ccgtcaatag atcagaccgt tttttattta ccattcatca attatattcc 1440
gcagtgggtt ggggtaaaaa aaatagaaga aaaccgcagc ggaccaattc cataccgttt 1500
ttacatacaa ataaacatgg tgcgcaacgg tttattgtcc gcctcaaaaa tgaaatggac 1560
taaaccgcag ataaattaga ccgctttgtc cgctgcctcc attcatagac taaaaaaaaa 1620
caaccaaaaa aaaaatgggt ccacgcccac gattttacac gaggtttctt gtggcgtaag 1680
gacaaaactc aaaagttcat aacgtttggt cctaaccagg tgtaatggat taagtaacag 1740
tcaattttct tattatagct gtatccatta tgtccacata tgcattcata tacattacac 1800
tggttggtct aagtgtagtt agattacgaa gactttcaag ttccattttt tgggttaggag 1860
ataaacataa tttaatgata ccgacttttag cactctaggc tcaaaacaag tacagaagag 1920
aatagtttta tttcaaacct gttgcattgt tgtatcaatt aattgtgtta gtctttgtat 1980
attcttacat aacggtccaa gtttggttgaa atagtttact tactaaactt ttcctaattg 2040
ggtcaaattt tattttatag gaaaaggaga gattgcaact tgttccaagg acatgaactc 2100
ggatcttttc ttcgcggtgt taggaggttt gggatcaattc ggcattataa caagagccag 2160
aattaaactt gaagtagctc cgaaaagggt atgttaaatt tgtaaattat gcaactacag 2220
aaaattctat gaaatttatg aatgaacata tatgcatttt tggatttttg taggccaaagt 2280

```

ggtaaagggtt tctatacata gattttctccg aattcacaag agatcaagaa cgagtgatat 2340
 cgaaaacgga cgggtgtagat ttcttagaag gttccattat ggtggaccat ggcccaccgg 2400
 ataactggag atccacgtat tatccaccgt ccgatcactt gaggatcgcc tcaatgggtca 2460
 aacgacatcg tgatcatctac tgccttgaag tcgtcaagta ttacgacgaa acttctcaat 2520
 acacagtcaa cgaggtcctg acatacatat aatcataaat catacatgta taattgggag 2580
 atctttatgc attattcaat tatattaatt tacttttagtt atttaactta tgcaggaaat 2640
 ggaggagtta agcgatagtt taaacatgt aagagggttt atgtacgaga aagatgtgac 2700
 gtatatggat ttcctaaacc gagttcgaac cggagagcta aacctgaaat ccaaaggcca 2760
 atgggatgtt ccacatccat ggcttaatct cttcgtacca aaaactcaaa tctccaaatt 2820
 tgatgatggg gtttttaagg gtattatcct aagaaataac atcactagcg gtcctgttct 2880
 tgtttatcct atgaatcgca acaagtaagt ttaactcgat attgcaaaat ttactatcta 2940
 cattttcgtt ttggaatccg aaatattctt acaagctaatt tttatgcggc gtttttaggt 3000
 ggaatgatcg gatgtctgcc gctatacccg aggaagatgt attttatgcg gtaggggttt 3060
 taagatccgc gggttttgac aattggggagg cttttgatca agaaaacatg gaaatactga 3120
 agttttgtga ggatgctaatt atgggggtta tacaatatct tccttatcat tcatcacaag 3180
 aaggatgggt tagacatttt ggtccgaggt ggaatatatt cgtagagaga aaatataaat 3240
 atgatcccaa aatgatatta tcaccgggac aaaatatatt tcaaaaaata aactcgagtt 3300
 ag 3302

<210> 6

<211> 523

<212> PRT

<213> Arabidopsis thaliana

<400> 6

Met Ala Ser Tyr Asn Leu Arg Ser Gln Val Arg Leu Ile Ala Ile Thr
 1 5 10 15

Ile Val Ile Ile Ile Thr Leu Ser Thr Pro Ile Thr Thr Asn Thr Ser
 20 25 30

Pro Gln Pro Trp Asn Ile Leu Ser His Asn Glu Phe Ala Gly Lys Leu
 35 40 45

Thr Ser Ser Ser Ser Ser Val Glu Ser Ala Ala Thr Asp Phe Gly His
 50 55 60

Val Thr Lys Ile Phe Pro Ser Ala Val Leu Ile Pro Ser Ser Val Glu
 65 70 75 80

Asp Ile Thr Asp Leu Ile Lys Leu Ser Phe Asp Ser Gln Leu Ser Phe
 85 90 95

Pro Leu Ala Ala Arg Gly His Gly His Ser His Arg Gly Gln Ala Ser
 100 105 110

Ala Lys Asp Gly Val Val Val Asn Met Arg Ser Met Val Asn Arg Asp

115		120		125
Arg Gly Ile Lys Val Ser Arg Thr Cys Leu Tyr Val Asp Val Asp Ala				
130		135		140
Ala Trp Leu Trp Ile Glu Val Leu Asn Lys Thr Leu Glu Leu Gly Leu				
145		150		155
Thr Pro Val Ser Trp Thr Asp Tyr Leu Tyr Leu Thr Val Gly Gly Thr				
	165		170	175
Leu Ser Asn Gly Gly Ile Ser Gly Gln Thr Phe Arg Tyr Gly Pro Gln				
	180		185	190
Ile Thr Asn Val Leu Glu Met Asp Val Ile Thr Gly Lys Gly Glu Ile				
	195		200	205
Ala Thr Cys Ser Lys Asp Met Asn Ser Asp Leu Phe Phe Ala Val Leu				
	210		215	220
Gly Gly Leu Gly Gln Phe Gly Ile Ile Thr Arg Ala Arg Ile Lys Leu				
225		230		235
Glu Val Ala Pro Lys Arg Ala Lys Trp Leu Arg Phe Leu Tyr Ile Asp				
	245		250	255
Phe Ser Glu Phe Thr Arg Asp Gln Glu Arg Val Ile Ser Lys Thr Asp				
	260		265	270
Gly Val Asp Phe Leu Glu Gly Ser Ile Met Val Asp His Gly Pro Pro				
	275		280	285
Asp Asn Trp Arg Ser Thr Tyr Tyr Pro Pro Ser Asp His Leu Arg Ile				
	290		295	300
Ala Ser Met Val Lys Arg His Arg Val Ile Tyr Cys Leu Glu Val Val				
305		310		315
Lys Tyr Tyr Asp Glu Thr Ser Gln Tyr Thr Val Asn Glu Glu Met Glu				
	325		330	335
Glu Leu Ser Asp Ser Leu Asn His Val Arg Gly Phe Met Tyr Glu Lys				
	340		345	350
Asp Val Thr Tyr Met Asp Phe Leu Asn Arg Val Arg Thr Gly Glu Leu				
	355		360	365
Asn Leu Lys Ser Lys Gly Gln Trp Asp Val Pro His Pro Trp Leu Asn				

370	375	380
Leu Phe Val Pro Lys Thr Gln Ile Ser Lys Phe Asp Asp Gly Val Phe		
385	390	395 400
Lys Gly Ile Ile Leu Arg Asn Asn Ile Thr Ser Gly Pro Val Leu Val		
	405	410 415
Tyr Pro Met Asn Arg Asn Lys Trp Asn Asp Arg Met Ser Ala Ala Ile		
	420	425 430
Pro Glu Glu Asp Val Phe Tyr Ala Val Gly Phe Leu Arg Ser Ala Gly		
	435	440 445
Phe Asp Asn Trp Glu Ala Phe Asp Gln Glu Asn Met Glu Ile Leu Lys		
450	455	460
Phe Cys Glu Asp Ala Asn Met Gly Val Ile Gln Tyr Leu Pro Tyr His		
465	470	475 480
Ser Ser Gln Glu Gly Trp Val Arg His Phe Gly Pro Arg Trp Asn Ile		
	485	490 495
Phe Val Glu Arg Lys Tyr Lys Tyr Asp Pro Lys Met Ile Leu Ser Pro		
	500	505 510
Gly Gln Asn Ile Phe Gln Lys Ile Asn Ser Ser		
515	520	

<210> 7

<211> 2782

<212> DNA

<213> Arabidopsis thaliana

<400> 7

```

atgactaata ctctctgttt aagcctcatt accctaataa cgcttttttat aagtttaacc 60
ccaaccttaa tcaaatcaga tgagggcatt gatgttttct taccatatac actcaacctt 120
acggtcctaa ccgatccctt ctccatctct gccgcttctc acgacttcgg taacataacc 180
gacgaaaatc ccggcgccgt cctctgccct tcctccacca cggaggtggc tcgtctcttc 240
cgtttcgcta acggaggatt ctcttacaat aaagggtcaa ccagccccgc gtctactttc 300
aaagtggctg ctcgaggcca aggccactcc ctccgtggcc aagcctctgc acccgagggt 360
gtcgtcgtga acatgacgtg tctcgccatg gcggtctaac cagcggcggg tggtatctcg 420
gcagacggga cttacgctga cgtggctgcc gggacgatgt ggggtggatgt tctgaaggcg 480
gcggtggata gaggcgtctc gccggttaca tggacggatt atttgtatct cagcgtcggc 540
gggacgttgt cgaacgctgg aatcggtggt cagacgttta gacacggccc tcagattagt 600
aacgttcatt agcttgacgt tattaccggt acgtaaatac caaaacttca ctaatctcgt 660

```

```

tacaattttt taattttttg gtaatatataa ttttgtacgg ctcaactctt aattaagaat 720
gaaacagtat ctatgatctt ctagatgctc tttttttgtc tgcaagcttt aattgtagta 780
acatcagcga tatatatatc acatgcatgt gtattattga tgataatata taatgtttta 840
gttacaaatt tgattctcaa ggtaaaactc acacgccata accagtataa aactccaaaa 900
atcacgtttt ggtcagaaat acatatcctt cattaacagt agttatgcta taatttgtga 960
ttataaataa ctccggaggt tgttcacaat actaaatttc aggaaaaggt gaaatgatga 1020
cttgctctcc aaagttaaac cctgaattgt tctatggagt tttaggaggt ttgggtcaat 1080
tcggtattat aacgagggcc aggattgctg tggatcatgc acccacaagg gtatgtatca 1140
tgcactata gtgtaatcaa ttataattt taatgtagt gtcctaaatc caaaatttga 1200
tttgatttgg ttggaacgta cgtatatata ataagtcaaa aggctgattt tgaagacgaa 1260
tttatatact tttgttgaat taaatctgat tttgcttacg ttttattaga ttctgcgtaa 1320
taaactctag gacttgctcg agtgaatct tgtcttatgc ttgcaaactc tgttgatgct 1380
aatatcta atcttttttatt atatttcctt acgtaagtt tagatatagt tattttaaac 1440
tgctataaat tgtgtacgta tagacttttag ataaaaaggt gtggtcgtt gcacctattt 1500
gtttatcgct atagtgatc aaaggctct atagtattct tgggttttct ttttggaaaa 1560
aatagaccat acaatccaag gaagatgatc ttaaatggac taatttatgg atataaattg 1620
atatacaaat ctgcaggtga aatggctctc cactactctac agtgacttct cggcttttaa 1680
aagagaccaa gagcgtttta tatcaatgac caatgatctc ggagttgact ttttggagg 1740
tcaacttatg atgtcaaatg gcttcgtaga cactctttt tcccactct cggatcaaac 1800
aagagtcgca tctcttgtga atgaccacc gatcatctat gttctcgaag tagccaagta 1860
ttatgacaga accaccctt ccattattga ccaggacta aaatccatta ttcattgatga 1920
ttatcttcac acaatcagta tcatcaccaa ttaccatcat cacttgctat atatgatcca 1980
aagtaaatat atcacatgat ataaataaat cgttcaaact ttttttttta aagaataaaa 2040
gaatcatttt caagcattac tcatacacat ctacgaatca ccgtgaccat atataaccat 2100
acgcttatta aataatcatt tttgtttgta ggtgattgac acgttaagta gaactctagg 2160
tttcgctcca gggtttatgt tcgtacaaga tgttcctgat ttcgatttct tgaaccgtgt 2220
ccgaaacgaa gaagataaac tcagatcttt aggactatgg gaagttcctc atccatggct 2280
taacatcttt gtcccggggt ctcgaatcca agattttcat gatggtgta ttaatggcct 2340
tcttctaaac caaacctcaa cttctgggtg tactctcttc tatcccacaa accgaaacaa 2400
gtaaataatt actttttgat tttgttttat ttgaaagtat atcccaataa tgtatgttaa 2460
attgttaaca agaatttatt ttattaatag atggaacaac cgcattgtcaa cgatgacacc 2520
ggacgaagat gttttttatg tgatcggatt actgcaatca gctgggtgat ctcaaaattg 2580
gcaagaactt gaaaatctca acgacaaggt tattcagttt tgtgaaaact cgggaattaa 2640
gattaaggaa tatttgatgc actatacaag aaaagaagat tgggttaaac attttgacc 2700
aaaatgggat gattttttta gaaagaaaat tatgtttgat cccaaaagac tattgtctcc 2760
aggacaagac atatttaatt aa 2782

```

<210> 8

<211> 524

<212> PRT

<213> Arabidopsis thaliana

<400> 8

Met Thr Asn Thr Leu Cys Leu Ser Leu Ile Thr Leu Ile Thr Leu Phe

1

5

10

15

Ile Ser Leu Thr Pro Thr Leu Ile Lys Ser Asp Glu Gly Ile Asp Val

20	25	30
Phe Leu Pro Ile Ser Leu Asn Leu Thr Val Leu Thr Asp Pro Phe Ser		
35	40	45
Ile Ser Ala Ala Ser His Asp Phe Gly Asn Ile Thr Asp Glu Asn Pro		
50	55	60
Gly Ala Val Leu Cys Pro Ser Ser Thr Thr Glu Val Ala Arg Leu Leu		
65	70	75
80		
Arg Phe Ala Asn Gly Gly Phe Ser Tyr Asn Lys Gly Ser Thr Ser Pro		
85	90	95
Ala Ser Thr Phe Lys Val Ala Ala Arg Gly Gln Gly His Ser Leu Arg		
100	105	110
Gly Gln Ala Ser Ala Pro Gly Gly Val Val Val Asn Met Thr Cys Leu		
115	120	125
Ala Met Ala Ala Lys Pro Ala Ala Val Val Ile Ser Ala Asp Gly Thr		
130	135	140
Tyr Ala Asp Val Ala Ala Gly Thr Met Trp Val Asp Val Leu Lys Ala		
145	150	155
160		
Ala Val Asp Arg Gly Val Ser Pro Val Thr Trp Thr Asp Tyr Leu Tyr		
165	170	175
Leu Ser Val Gly Gly Thr Leu Ser Asn Ala Gly Ile Gly Gly Gln Thr		
180	185	190
Phe Arg His Gly Pro Gln Ile Ser Asn Val His Glu Leu Asp Val Ile		
195	200	205
Thr Gly Lys Gly Glu Met Met Thr Cys Ser Pro Lys Leu Asn Pro Glu		
210	215	220
Leu Phe Tyr Gly Val Leu Gly Gly Leu Gly Gln Phe Gly Ile Ile Thr		
225	230	235
240		
Arg Ala Arg Ile Ala Leu Asp His Ala Pro Thr Arg Val Lys Trp Ser		
245	250	255
Arg Ile Leu Tyr Ser Asp Phe Ser Ala Phe Lys Arg Asp Gln Glu Arg		
260	265	270
Leu Ile Ser Met Thr Asn Asp Leu Gly Val Asp Phe Leu Glu Gly Gln		

275	280	285
Leu Met Met Ser Asn Gly Phe Val Asp Thr Ser Phe Phe Pro Leu Ser		
290	295	300
Asp Gln Thr Arg Val Ala Ser Leu Val Asn Asp His Arg Ile Ile Tyr		
305	310	315 320
Val Leu Glu Val Ala Lys Tyr Tyr Asp Arg Thr Thr Leu Pro Ile Ile		
	325	330 335
Asp Gln Val Ile Asp Thr Leu Ser Arg Thr Leu Gly Phe Ala Pro Gly		
	340	345 350
Phe Met Phe Val Gln Asp Val Pro Tyr Phe Asp Phe Leu Asn Arg Val		
	355	360 365
Arg Asn Glu Glu Asp Lys Leu Arg Ser Leu Gly Leu Trp Glu Val Pro		
	370	375 380
His Pro Trp Leu Asn Ile Phe Val Pro Gly Ser Arg Ile Gln Asp Phe		
385	390	395 400
His Asp Gly Val Ile Asn Gly Leu Leu Leu Asn Gln Thr Ser Thr Ser		
	405	410 415
Gly Val Thr Leu Phe Tyr Pro Thr Asn Arg Asn Lys Trp Asn Asn Arg		
	420	425 430
Met Ser Thr Met Thr Pro Asp Glu Asp Val Phe Tyr Val Ile Gly Leu		
	435	440 445
Leu Gln Ser Ala Gly Gly Ser Gln Asn Trp Gln Glu Leu Glu Asn Leu		
	450	455 460
Asn Asp Lys Val Ile Gln Phe Cys Glu Asn Ser Gly Ile Lys Ile Lys		
465	470	475 480
Glu Tyr Leu Met His Tyr Thr Arg Lys Glu Asp Trp Val Lys His Phe		
	485	490 495
Gly Pro Lys Trp Asp Asp Phe Leu Arg Lys Lys Ile Met Phe Asp Pro		
	500	505 510
Lys Arg Leu Leu Ser Pro Gly Gln Asp Ile Phe Asn		
	515	520

<210> 9
 <211> 2805
 <212> DNA
 <213> Arabidopsis thaliana

<400> 9
 atgacgtcaa gctttcttct cctgacgttc gccatatgta aactgatcat agccgtgggt 60
 ctaaacgtgg gccccagtga gctcctccgc atcggagcca tagatgtcga cggccacttc 120
 accgtccacc cttccgactt agcctccgtc tcctcagact tcggtatgct gaagtcacct 180
 gaagagccat tggcgtgct tcatccatca tcggccgaag acgtggcacg actcgtcaga 240
 acagcttacg gttcagccac ggcgtttccg gtctcagccc gaggccacgg ccattccata 300
 aacggacaag ccgcggcggg gaggaacggt gtggtgggtt aaatgaacca cggcgtaacc 360
 gggacgccc agccactcgt ccgaccggat gaaatgtatg tggatgtatg ggggtggagag 420
 ttatgggtcg atgtgttgaa gaaaacgttg gagcatggct tagcaccaaa atcatggacg 480
 gattacttgt atctaaccgt tggaggtaca ctctccaatg caggaatcag tggtaagct 540
 tttcaccatg gtctcaaat tagtaacgtc cttgagctcg acgttgtaac tggtagtat 600
 taaaacattc aagttcatat attttaaatg cttttgtctg aagttttact aataacaaga 660
 aattgatacc aaaaagtagg gaaaggagag gtgatgagat gtcagaaga agagaacaca 720
 aggctattcc atggagtctt tgggtggatta ggtcaatttg ggatcatcac tcgagcacga 780
 atctctctcg aaccagctcc ccaaagggtg atattttttt aatgactagc tatcaaaaat 840
 ccctggcggg tccatacgtt gtaatctttt tagtttttac tgttgatggt attttttata 900
 tattttggat aataaaaccc taaaatggta tattgtgatg acaggtgaga tggatacggg 960
 tattgtatcc gagcttcaaa gtgtttacgg aggaccaaga gtacttaatc tcaatgcatg 1020
 gtcaattaaa gtttgattac gtggaaggtt ttgtgattgt ggacgaagga ctctgtaaca 1080
 attggagatc ttctttcttc tctccacgta acccgtcaa gatctcctct gttagttcca 1140
 acggctctgt tttgtattgc cttgagatca ccaagaacta ccacgactcc gactccgaaa 1200
 tcgttgatca ggtcactttc attattcact tagaaaaaag cgatattttc attttttata 1260
 ttgatgaata tctggaagga ttaacgcta tgcgactatt gggaaatcat tatgaaaaaa 1320
 tatttagttt atatgattga aagtggcttc catagtattt ttgttggtgc gactttatta 1380
 taacttaaat ttggaagagg acatgaagaa gaagccagag aggatctaca gagatctagc 1440
 ttttccacct gaacttaata atgcacattt atataattat ttttcttctt ctaaagttta 1500
 gtttatcact agcgaattaa tcatgggtac taattaagta gtggacaggg tcatggacca 1560
 ctcaactcacc aaataatgat tcctctttac tcttaagttt aattttaata aaaccaactc 1620
 tactggaatc ttaacttata cttgggtttt gtaggctttt atagcaacac ggttttttta 1680
 attttcctat tccagatttt gtatattaaa tgtcgatttt ttttcttttt gtttcaggaa 1740
 gttgagattc tgatgaagaa attgaatttc ataccgacat cggctctttac aacggattta 1800
 caatatgtgg actttctcga ccgggtacac aaggccgaat tgaagctccg gtccaagaat 1860
 ttatgggagg ttccacaccc atggctcaac ctcttcgtgc caaaatcaag aatctctgac 1920
 ttcgataaag gcgttttcaa gggcattttg ggaaataaaa caagtggccc tattcttata 1980
 taccatga acaaagacaa gtaagtcttg acattaccat tgattactac ttctaaattt 2040
 cttctctaga aaaaagaata aaacgagttt tgcattgcat gcatgcaaag ttacacttgt 2100
 ggggattaat tagtgggtcca agaaaaaag tttgtcaaaa ttgaaaaaaa ctagacacgt 2160
 ggtacatggg attgtccgaa aaacgttgct cacatgtgca tcgaaccagc taagattgac 2220
 aacaacactt cgtcggctcg tatttctctt tttgttttgt gaccaaatec gatgggtccag 2280
 attgggttta tttgttttta agttcctaga actcatggtg ggtgggtccc aatcagattc 2340
 tcttagacca aaccgatctc aacgaacctt ccgcacatca ttgattatta cattaatata 2400
 gatattgtcg ttgctgacgt gtcgtaattt gatgttattg tcagatggga cgagaggagc 2460

tcagccgtga cgccggatga ggaagttttc tatctggtgg ctctattgag atcagcttta 2520
acggacggtg aagagacaca gaagctagag tatctgaaag atcagaaccg tcggatcttg 2580
gagttctgtg aacaagccaa gatcaatgtg aagcagtatc ttcctcacca cgcaacacag 2640
gaagagtggg tggctcattt tggggacaag tgggatcggg tcagaagctt aaaggctgag 2700
tttgatccgc gacacatact cgctactggg cagagaatct ttcaaaaccc atctttgtct 2760
ttgtttcctc cgtcgtcgtc ttcttcgtca gcggcttcat ggtga 2805

<210> 10

<211> 536

<212> PRT

<213> Arabidopsis thaliana

<400> 10

Met Thr Ser Ser Phe Leu Leu Leu Thr Phe Ala Ile Cys Lys Leu Ile
1 5 10 15

Ile Ala Val Gly Leu Asn Val Gly Pro Ser Glu Leu Leu Arg Ile Gly
20 25 30

Ala Ile Asp Val Asp Gly His Phe Thr Val His Pro Ser Asp Leu Ala
35 40 45

Ser Val Ser Ser Asp Phe Gly Met Leu Lys Ser Pro Glu Glu Pro Leu
50 55 60

Ala Val Leu His Pro Ser Ser Ala Glu Asp Val Ala Arg Leu Val Arg
65 70 75 80

Thr Ala Tyr Gly Ser Ala Thr Ala Phe Pro Val Ser Ala Arg Gly His
85 90 95

Gly His Ser Ile Asn Gly Gln Ala Ala Ala Gly Arg Asn Gly Val Val
100 105 110

Val Glu Met Asn His Gly Val Thr Gly Thr Pro Lys Pro Leu Val Arg
115 120 125

Pro Asp Glu Met Tyr Val Asp Val Trp Gly Gly Glu Leu Trp Val Asp
130 135 140

Val Leu Lys Lys Thr Leu Glu His Gly Leu Ala Pro Lys Ser Trp Thr
145 150 155 160

Asp Tyr Leu Tyr Leu Thr Val Gly Gly Thr Leu Ser Asn Ala Gly Ile
165 170 175

Ser Gly Gln Ala Phe His His Gly Pro Gln Ile Ser Asn Val Leu Glu

180	185	190
Leu Asp Val Val Thr Gly Lys Gly Glu Val Met Arg Cys Ser Glu Glu 195 200 205		
Glu Asn Thr Arg Leu Phe His Gly Val Leu Gly Gly Leu Gly Gln Phe 210 215 220		
Gly Ile Ile Thr Arg Ala Arg Ile Ser Leu Glu Pro Ala Pro Gln Arg 225 230 235 240		
Val Arg Trp Ile Arg Val Leu Tyr Ser Ser Phe Lys Val Phe Thr Glu 245 250 255		
Asp Gln Glu Tyr Leu Ile Ser Met His Gly Gln Leu Lys Phe Asp Tyr 260 265 270		
Val Glu Gly Phe Val Ile Val Asp Glu Gly Leu Val Asn Asn Trp Arg 275 280 285		
Ser Ser Phe Phe Ser Pro Arg Asn Pro Val Lys Ile Ser Ser Val Ser 290 295 300		
Ser Asn Gly Ser Val Leu Tyr Cys Leu Glu Ile Thr Lys Asn Tyr His 305 310 315 320		
Asp Ser Asp Ser Glu Ile Val Asp Gln Glu Val Glu Ile Leu Met Lys 325 330 335		
Lys Leu Asn Phe Ile Pro Thr Ser Val Phe Thr Thr Asp Leu Gln Tyr 340 345 350		
Val Asp Phe Leu Asp Arg Val His Lys Ala Glu Leu Lys Leu Arg Ser 355 360 365		
Lys Asn Leu Trp Glu Val Pro His Pro Trp Leu Asn Leu Phe Val Pro 370 375 380		
Lys Ser Arg Ile Ser Asp Phe Asp Lys Gly Val Phe Lys Gly Ile Leu 385 390 395 400		
Gly Asn Lys Thr Ser Gly Pro Ile Leu Ile Tyr Pro Met Asn Lys Asp 405 410 415		
Lys Trp Asp Glu Arg Ser Ser Ala Val Thr Pro Asp Glu Glu Val Phe 420 425 430		
Tyr Leu Val Ala Leu Leu Arg Ser Ala Leu Thr Asp Gly Glu Glu Thr		

435	440	445
Gln Lys Leu Glu Tyr Leu Lys Asp Gln Asn Arg Arg Ile Leu Glu Phe		
450	455	460
Cys Glu Gln Ala Lys Ile Asn Val Lys Gln Tyr Leu Pro His His Ala		
465	470	475 480
Thr Gln Glu Glu Trp Val Ala His Phe Gly Asp Lys Trp Asp Arg Phe		
485	490	495
Arg Ser Leu Lys Ala Glu Phe Asp Pro Arg His Ile Leu Ala Thr Gly		
500	505	510
Gln Arg Ile Phe Gln Asn Pro Ser Leu Ser Leu Phe Pro Pro Ser Ser		
515	520	525
Ser Ser Ser Ser Ala Ala Ser Trp		
530	535	

<210> 11
 <211> 1936
 <212> DNA
 <213> Arabidopsis thaliana

<400> 11

atgcttatag taagaagttt caccatcttg cttctcagct gcatagcctt taagttggct	60
tgctgcttct ctagcagcat ttcttctttg aaggcgcttc ccctagtagg ccatttggag	120
tttgaacatg tccatcacgc ctccaaagat tttggaaatc gataccagtt gatccctttg	180
gcggtcttac atcccaaate ggtaagcgac atcgctcaa cgatacgaca catctggatg	240
atggggcactc attcacagct tacagtggca gcgagaggtc gtggacattc actccaaggc	300
caagctcaaa caagacatgg aattgttata cacatggaat cactccatcc ccagaagctg	360
caggtctaca gtgtggattc ccctgctcca tatgttgatg tgtctggtgg tgagctgtgg	420
ataaacattt tgcattgagac cctcaagtac gggcttgac caaatcatg gacggattac	480
ctgcatttaa ctgtagggtg tactctgtcc aatgctggaa taagcggcca ggcattccga	540
catggaccac agatcagcaa tgttcatcaa ctggagattg tcacaggtta gttcagagtt	600
gcagtattcg tgttttgaaa gcatagactc tatatggttg gtgactatta acaacatgaa	660
gagattcccg agaatagcta cccactaatg tcatgcctat ttattgactg caggaaaagg	720
cgagatccta aactgtacaa agaggcagaa cagcgactta tttaatggtg ttcttggtgg	780
tttaggtcag tttggcatca taacgcgggc aagaatagca ttggaaccag caccaaccat	840
ggtaaacaat aaataaataa aaaacttaaa aactgaacac gcgtgtgtcc tcctaactct	900
gtataatgga caggtaaaat ggataagagt gttatacctg gattttgcag cttttgccaa	960
ggaccaagag caactaatat ctgcccaggg ccacaaattc gattacatag aagggtttgt	1020
gataataaac aggacaggcc tcctgaacag ctggaggttg tctttcaccg cagaagagcc	1080
tttagaagca agccaattca agtttgatgg aaggactctg tattgtctgg agctagccaa	1140
gtatttgaag caagataaca aagacgtaat caaccaggtg agaaaacaga gtagaagcaa	1200

tcggtagaat cttctttggt agatgacatt cattggaact gaaaatatat atatatttgt 1260
 ccaatccagg aagtgaaga aacattatca gagctaagct acgtgacgtc gacactgttt 1320
 acaacggagg tagcatatga agcattcttg gacagggtac atgtgtctga ggtaaaactc 1380
 cgatcgaaag ggcagtggga ggtgccacat ccatggctga acctcctggt accaagaagc 1440
 aaaatcaatg aatttgcaag aggtgtattt ggaaacatac taacggatac aagcaacggc 1500
 ccagtcatcg tctaccagc gaacaaatca aagtaagaaa gaaagaaaga aagagctagt 1560
 catgattttg tttcttttca cttgttgaca aaacaaaagc atgttggtga gcagggtggga 1620
 caatcaaaca tcagcagtaa caccggagga agaggtattc tacctggtgg cgatcctaac 1680
 atcggcattc ccagggtcgg caggaaagga tggagtagaa gagatcttga ggcggaacag 1740
 aagaatactg gaattcagtg aagaagcagg gatagggttg aagcagtatc tgccacatta 1800
 cagcacaaga gaagagtggga gatcccatct cgggggacaag tggggagaat ttgtgaggag 1860
 gaaatccaga tatgatccat tggcaattct tgcgcctggc caccgaattt ttcaaaaggc 1920
 agtctcatat tcatga 1936

<210> 12

<211> 504

<212> PRT

<213> Arabidopsis thaliana

<400> 12

Met	Leu	Ile	Val	Arg	Ser	Phe	Thr	Ile	Leu	Leu	Leu	Ser	Cys	Ile	Ala
1				5					10					15	
Phe	Lys	Leu	Ala	Cys	Cys	Phe	Ser	Ser	Ser	Ile	Ser	Ser	Leu	Lys	Ala
			20					25					30		
Leu	Pro	Leu	Val	Gly	His	Leu	Glu	Phe	Glu	His	Val	His	His	Ala	Ser
		35					40					45			
Lys	Asp	Phe	Gly	Asn	Arg	Tyr	Gln	Leu	Ile	Pro	Leu	Ala	Val	Leu	His
	50					55					60				
Pro	Lys	Ser	Val	Ser	Asp	Ile	Ala	Ser	Thr	Ile	Arg	His	Ile	Trp	Met
	65				70					75					80
Met	Gly	Thr	His	Ser	Gln	Leu	Thr	Val	Ala	Ala	Arg	Gly	Arg	Gly	His
			85						90					95	
Ser	Leu	Gln	Gly	Gln	Ala	Gln	Thr	Arg	His	Gly	Ile	Val	Ile	His	Met
		100						105					110		
Glu	Ser	Leu	His	Pro	Gln	Lys	Leu	Gln	Val	Tyr	Ser	Val	Asp	Ser	Pro
		115					120					125			
Ala	Pro	Tyr	Val	Asp	Val	Ser	Gly	Gly	Glu	Leu	Trp	Ile	Asn	Ile	Leu
	130						135					140			

His	Glu	Thr	Leu	Lys	Tyr	Gly	Leu	Ala	Pro	Lys	Ser	Trp	Thr	Asp	Tyr		
145					150					155					160		
Leu	His	Leu	Thr	Val	Gly	Gly	Thr	Leu	Ser	Asn	Ala	Gly	Ile	Ser	Gly		
				165					170					175			
Gln	Ala	Phe	Arg	His	Gly	Pro	Gln	Ile	Ser	Asn	Val	His	Gln	Leu	Glu		
			180					185					190				
Ile	Val	Thr	Gly	Lys	Gly	Glu	Ile	Leu	Asn	Cys	Thr	Lys	Arg	Gln	Asn		
			195				200					205					
Ser	Asp	Leu	Phe	Asn	Gly	Val	Leu	Gly	Gly	Leu	Gly	Gln	Phe	Gly	Ile		
	210					215					220						
Ile	Thr	Arg	Ala	Arg	Ile	Ala	Leu	Glu	Pro	Ala	Pro	Thr	Met	Asp	Gln		
225					230					235					240		
Glu	Gln	Leu	Ile	Ser	Ala	Gln	Gly	His	Lys	Phe	Asp	Tyr	Ile	Glu	Gly		
				245					250					255			
Phe	Val	Ile	Ile	Asn	Arg	Thr	Gly	Leu	Leu	Asn	Ser	Trp	Arg	Leu	Ser		
			260					265					270				
Phe	Thr	Ala	Glu	Glu	Pro	Leu	Glu	Ala	Ser	Gln	Phe	Lys	Phe	Asp	Gly		
		275					280					285					
Arg	Thr	Leu	Tyr	Cys	Leu	Glu	Leu	Ala	Lys	Tyr	Leu	Lys	Gln	Asp	Asn		
	290					295					300						
Lys	Asp	Val	Ile	Asn	Gln	Glu	Val	Lys	Glu	Thr	Leu	Ser	Glu	Leu	Ser		
305				310					315					320			
Tyr	Val	Thr	Ser	Thr	Leu	Phe	Thr	Thr	Glu	Val	Ala	Tyr	Glu	Ala	Phe		
				325					330					335			
Leu	Asp	Arg	Val	His	Val	Ser	Glu	Val	Lys	Leu	Arg	Ser	Lys	Gly	Gln		
			340					345					350				
Trp	Glu	Val	Pro	His	Pro	Trp	Leu	Asn	Leu	Leu	Val	Pro	Arg	Ser	Lys		
		355					360					365					
Ile	Asn	Glu	Phe	Ala	Arg	Gly	Val	Phe	Gly	Asn	Ile	Leu	Thr	Asp	Thr		
	370					375					380						
Ser	Asn	Gly	Pro	Val	Ile	Val	Tyr	Pro	Val	Asn	Lys	Ser	Lys	Trp	Asp		
385					390					395					400		

Asn Gln Thr Ser Ala Val Thr Pro Glu Glu Glu Val Phe Tyr Leu Val
 405 410 415
 Ala Ile Leu Thr Ser Ala Ser Pro Gly Ser Ala Gly Lys Asp Gly Val
 420 425 430
 Glu Glu Ile Leu Arg Arg Asn Arg Arg Ile Leu Glu Phe Ser Glu Glu
 435 440 445
 Ala Gly Ile Gly Leu Lys Gln Tyr Leu Pro His Tyr Thr Thr Arg Glu
 450 455 460
 Glu Trp Arg Ser His Phe Gly Asp Lys Trp Gly Glu Phe Val Arg Arg
 465 470 475 480
 Lys Ser Arg Tyr Asp Pro Leu Ala Ile Leu Ala Pro Gly His Arg Ile
 485 490 495
 Phe Gln Lys Ala Val Ser Tyr Ser
 500

<210> 13
 <211> 31
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence:oligonucleotide

<400> 13
 cggctcgacat gggattgacc tcataccttac g

31

<210> 14
 <211> 35
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence:oligonucleotide

<400> 14
 gcgctcgactt atacagttct aggtttcggc agtat

35

<210> 15

<211> 33
 <212> DNA
 <213> Artificial Sequence

 <220>
 <223> Description of Artificial Sequence:oligonucleotide

 <400> 15
 gcggtaccag agagagaaac ataaacaaat ggc 33

 <210> 16
 <211> 31
 <212> DNA
 <213> Artificial Sequence

 <220>
 <223> Description of Artificial Sequence:oligonucleotide

 <400> 16
 gcggtaccga attttacttc caccaaaatg c 31

 <210> 17
 <211> 34
 <212> DNA
 <213> Artificial Sequence

 <220>
 <223> Description of Artificial Sequence:oligonucleotide

 <400> 17
 gcggtacctt cattgataag aatcaagcta ttca 34

 <210> 18
 <211> 31
 <212> DNA
 <213> Artificial Sequence

 <220>
 <223> Description of Artificial Sequence:oligonucleotide

 <400> 18
 gcggtaccga aagtggtag aacgactaac a 31

 <210> 19

<211> 28
<212> DNA
<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:oligonucleotide

<400> 19

gcggtacccc cattaaccta cccgtttg

28

<210> 20
<211> 32
<212> DNA
<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:oligonucleotide

<400> 20

gcggtaccag acgatgaacg tacttgtctg ta

32

<210> 21
<211> 31
<212> DNA
<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:oligonucleotide

<400> 21

ggggtaccgt gaaatgacgt caagctttct t

31

<210> 22
<211> 31
<212> DNA
<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:oligonucleotide

<400> 22

ggggtaccct ttcctcttgg ttttgcctg t

31

<210> 23

<211> 32
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:oligonucleotide

<400> 23
gctctagatc aggaaaagaa ccatgcttat ag 32

<210> 24
<211> 32
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:oligonucleotide

<400> 24
gctctagatc atgagtatga gactgccttt tg 32